



www.combinatorics.ir

---

Transactions on Combinatorics

ISSN (print): 2251-8657, ISSN (on-line): 2251-8665

Vol. 7 No. 4 (2018), pp. 7-10.

© 2018 University of Isfahan

---



www.ui.ac.ir

## A NOTE ON 1-FACTORIZABILITY OF QUARTIC SUPERSOLVABLE CAYLEY GRAPHS

MILAD AHANJIDEH AND ALI IRANMANESH\*

Communicated by Tommy R. Jensen

ABSTRACT. Alspach et al. conjectured that every quartic Cayley graph on an even solvable group is 1-factorizable. In this paper, we verify this conjecture for quartic Cayley graphs on supersolvable groups of even order.

### 1. Introduction and Preliminary Results

Let  $G$  be a finite group with identity 1 and  $S \subset G \setminus \{1\}$ . A *Cayley graph* with respect to the set  $S$ , denoted by  $\text{Cay}(G, S)$ , is a graph whose vertex set is the set of elements of  $G$  with adjacency defined by

$$g \sim h \text{ if and only if } g^{-1}h \in S \cup S^{-1}$$

for every  $g, h \in G$ , where  $S^{-1} = \{s^{-1} \mid s \in S\}$ . We see at once that if  $S$  generates  $G$ , then  $\text{Cay}(G, S)$  is connected.

A group  $G$  is *supersolvable* if there exists a normal series

$$\{1\} = H_0 \triangleleft H_1 \triangleleft \cdots \triangleleft H_n = G$$

such that each quotient group  $H_i/H_{i-1}$  is cyclic. Note that every supersolvable group is a solvable group.

---

MSC(2010): Primary: 05C25; Secondary: 05C70.

Keywords: Cayley graph, 1-factorization, supersolvable group.

Received: 04 June 2017, Accepted: 29 May 2018.

\*Corresponding author.

DOI: <http://dx.doi.org/10.22108/toc.2018.104578.1500>

A  $j$ -factor of a graph is a spanning subgraph which is regular of valence  $j$ . In particular, a 1-factor of a graph is a collection of edges such that each vertex is incident with exactly one edge. A 1-factorization of a regular graph is a partition of the edge set of the graph into disjoint 1-factors. A 1-factorization of a regular graph of valence  $v$  is equivalent to a coloring of the edges in  $v$  colors (coloring each 1-factor a different color).

Stong investigated about 1-factorizability of Cayley graphs in 1985 [4]. After that, Alspach et al. [3] studied the factorization of quartic Cayley graphs on some solvable groups of even order. They posed the following conjecture:

**Conjecture 1.1.** [3] *Every quartic Cayley graph on an even solvable group is 1-factorizable.*

They proved their conjecture for an even solvable group  $G$  such that the commutator subgroup  $G'$  is an elementary abelian  $p$ -group. Also Abdollahi showed that every Cayley graph on a nilpotent group of even order is 1-factorizable [1].

In this paper, we verify Conjecture 1.1 for an even supersolvable group. In fact we prove the following theorem:

**Theorem 1.2.** *Every quartic Cayley graph on an even supersolvable group is 1-factorizable*

To prove the above theorem, we need several lemmas.

**Lemma 1.3.** [2] *Every cubic Cayley graph on a solvable group is 3-edge-colorable.*

The following lemmas are due to Stong [4].

**Lemma 1.4.** *Let  $S_1, S_2 \subseteq G \setminus \{1\}$ , not necessarily generating sets. Suppose  $\text{Cay}(G, S_2)$  is 1-factorizable and  $S_2 \subseteq S_1$ . If every element in  $S_1 \setminus S_2$  has even order then  $\text{Cay}(G, S_1)$  is 1-factorizable.*

**Lemma 1.5.** *If  $G$  is a 2-generated group of even order with a cyclic commutator subgroup, then  $\text{Cay}(G, \{a, a^{-1}, b, b^{-1}\})$  is 1-factorizable*

**Lemma 1.6.** *Suppose that  $N$  is a normal subgroup of  $G$  and  $S$  is a generating set of  $G$  disjoint from  $N$ . Assume that when  $s_i \neq s_j^{\pm 1}$ , neither  $s_i s_j$  nor  $s_i s_j^{-1}$  belongs to  $N$ . If  $\text{Cay}(G/N, S/N)$  is 1-factorizable, then so is  $\text{Cay}(G, S)$ .*

## 2. The proof of Theorem 1.2

The proof falls naturally into three parts. First, assume that all elements of  $S$  have order 2. Since the edges generated by an element of order 2 form a 1-factor, it follows that  $\text{Cay}(G, S)$  is 1-factorizable. If  $S = \{a, a^{-1}, b, c\}$ , where  $O(a) > 2$  and  $O(b) = O(c) = 2$ , then by Lemma 1.3,  $\text{Cay}(G, S \setminus \{c\})$  is 1-factorizable and hence, Lemma 1.4 completes the proof. Now, assume that  $S = \{a, a^{-1}, b, b^{-1}\}$  where  $O(a), O(b) > 2$  and let  $N$  be a minimal normal subgroup of  $G$ . Suppose that the theorem is false and let  $G$  be the smallest supersolvable group in the question such that  $\text{Cay}(G, S)$  is not

1-factorizable. Suppose that  $\langle S \rangle \neq G$ . Since  $\langle S \rangle$  is a proper supersolvable subgroup of  $G$ , we can see that  $\text{Cay}(\langle S \rangle, S)$  is 1-factorizable by our assumption. Now let  $T = \{x_1, \dots, x_t\}$ , where  $t \in \mathbb{N}$ , be a left transversal set of  $\langle S \rangle$  in  $G$ . Thus  $\{x_i \text{Cay}(\langle S \rangle, S) : 1 \leq i \leq t\}$  is the set of the connected components of  $\text{Cay}(G, S)$  where for every  $1 \leq i \leq t$ ,  $x_i \text{Cay}(\langle S \rangle, S)$  is a graph which its vertex set is  $x_i \langle S \rangle$  and two vertices  $x_i y_j$  and  $x_i y_k$  are adjacent if and only if  $(x_i y_j)^{-1} (x_i y_k) \in S$ . Therefore for every  $1 \leq i \leq t$ ,  $x_i \text{Cay}(\langle S \rangle, S)$  and  $\text{Cay}(\langle S \rangle, S)$  are isomorphic and hence  $x_i \text{Cay}(\langle S \rangle, S)$  is 1-factorizable. So  $\text{Cay}(G, S)$  is 1-factorizable which is a contradiction. Thus let  $\langle S \rangle = G$ . We continue the proof in two cases:

**Case1.** Suppose that  $N \cap S \neq \emptyset$ . If  $a, b \in N$ , then  $N = G$ . Moreover,  $|G|$  is even and  $N$  is a cyclic group of prime order. So,  $N = G = \mathbb{Z}_2$  and the proof is complete. If  $a \in N$ , then  $G/N = \langle bN \rangle$  is abelian and hence,  $G' \leq N$ . From this we have  $G'$  is cyclic. Lemma 1.5 shows  $\text{Cay}(G, S)$  is 1-factorizable. This is a contradiction.

**Case2.** Let  $N \cap S = \emptyset$ . The proof will be divided into two subcases.

**Subcase (a).** Let  $|N|$  be odd. Since  $|N|$  is odd and  $|G|$  is even,  $|G/N|$  is even. If  $aN \in \{bN, b^{-1}N\}$ , then  $G/N = \langle aN, bN \rangle = \langle aN \rangle$ . So  $O(aN) = O(bN)$  is even and therefore  $O(a), O(b)$  are even. Thus  $\text{Cay}(G, \{a, a^{-1}\})$  is a union of cycles of even lengths which is 1-factorizable. Lemma 1.4 shows that  $\text{Cay}(G, S)$  is 1-factorizable. This is impossible. Let  $aN \notin \{bN, b^{-1}N\}$ . Since  $|S| = |S/N|$ , we conclude that  $\text{Cay}(G/N, S/N)$  is 1-factorizable by our assumption, and so is  $\text{Cay}(G, S)$  from Lemma 1.6, a contradiction.

**Subcase (b).** Suppose that  $|N|$  is even. So  $|N| = 2$ . If  $|G/N|$  is even, then the same argument as used in Subcase (a) shows that  $\text{Cay}(G, S)$  is 1-factorizable. This is a contradiction. Now, let  $|G/N| = m$  be odd. Since  $|N| = 2$ ,  $N \leq Z(G)$ , and hence we have  $G = M \times N$ , where  $M$  is a Hall subgroup of order  $m$ . Note that  $G$  is a supersolvable group, so is  $M$ . Thus  $M$  has a minimal normal subgroup  $M_1$  of prime order  $p$ . Since  $G = M \times N$  and  $p \mid m$ , we get  $M_1$  is a minimal normal subgroup of  $G$  of odd order. Hence, by substituting  $N$  with  $M_1$  in Subcase (a), we see that  $\text{Cay}(G, S)$  is 1-factorizable, which is a contradiction. This completes the proof.

### Acknowledgments

The authors wish to thank to referee for reading the paper carefully and making the useful suggestions.

### REFERENCES

- [1] A. Abdollahi, 1-factorizations of Cayley graphs, *Ars Combin.*, **86** (2008) 129–131.
- [2] B. Alspach, Y. Liu and C. Zhang, Nowhere-zero 4-flows and Cayley graphs on solvable groups, *SIAM J. Discrete Math.*, **9** (1996) 151–154.
- [3] B. Alspach, M. Morton and Y. Qin, 1-factorizations of Cayley graphs on solvable groups, Report, The University of Auckland Library, <https://researchspace.auckland.ac.nz/bitstream/handle/2292/5039/393.pdf>.
- [4] R. Stong, On 1-factorizability of Cayley graphs, *J. Combin. Theory Ser. B*, **39** (1985) 298–307.

**Milad Ahanjideh**

Department of Mathematics, Faculty of Mathematical Sciences, Tarbiat Modares University, P.O.Box 14115-137, Tehran,  
Iran

Email: [ahanjideh@gmail.com](mailto:ahanjideh@gmail.com)

**Ali Iranmanesh**

Department of Mathematics, Faculty of Mathematical Sciences, Tarbiat Modares University, P.O.Box 14115-137, Tehran,  
Iran

Email: [iranmanesh@modares.ac.ir](mailto:iranmanesh@modares.ac.ir)