



www.combinatorics.ir

---

Transactions on Combinatorics

ISSN (print): 2251-8657, ISSN (on-line): 2251-8665

Vol. 01 No.1 (2012), pp. 7-13.

© 2012 University of Isfahan

---



www.ui.ac.ir

## **$k$ -TUPLE TOTAL DOMINATION AND MYCIELESKIAN GRAPHS**

A. P. KAZEMI

Communicated by Mehdi Alaeiyan

**ABSTRACT.** Let  $k$  be a positive integer. A subset  $S$  of  $V(G)$  in a graph  $G$  is a  $k$ -tuple total dominating set of  $G$  if every vertex of  $G$  has at least  $k$  neighbors in  $S$ . The  $k$ -tuple total domination number  $\gamma_{\times k,t}(G)$  of  $G$  is the minimum cardinality of a  $k$ -tuple total dominating set of  $G$ . In this paper for a given graph  $G$  with minimum degree at least  $k$ , we find some sharp lower and upper bounds on the  $k$ -tuple total domination number of the  $m$ -Mycieleskian graph  $\mu_m(G)$  of  $G$  in terms on  $k$  and  $\gamma_{\times k,t}(G)$ . Specially we give the sharp bounds  $\gamma_{\times k,t}(G) + 1$  and  $\gamma_{\times k,t}(G) + k$  for  $\gamma_{\times k,t}(\mu_1(G))$ , and characterize graphs with  $\gamma_{\times k,t}(\mu_1(G)) = \gamma_{\times k,t}(G) + 1$ .

### **1. Introduction**

In this paper,  $G = (V, E)$  is a simple graph with the *vertex set*  $V$  and the *edge set*  $E$ . The *order*  $|V|$  of  $G$  is denoted by  $n = n(G)$ . The *open neighborhood* and the *closed neighborhood* of a vertex  $v \in V$  are  $N_G(v) = \{u \in V(G) \mid uv \in E(G)\}$  and  $N_G[v] = N_G(v) \cup \{v\}$ , respectively. Also the *degree* of  $v$  is  $\deg_G(v) = |N_G(v)|$ . The *minimum* and *maximum degree* of  $G$  are denoted by  $\delta = \delta(G)$  and  $\Delta = \Delta(G)$ , respectively. We write  $K_n$  and  $C_n$  for the *complete graph* and the *cycle* of order  $n$ , respectively, while  $G[S]$  and  $K_{n_1, n_2, \dots, n_p}$  denote the *subgraph induced* on  $G$  by a vertex set  $S$ , and the *complete  $p$ -partite graph*, respectively.

Let  $S \subseteq V$  and let  $k$  be a positive integer. For each  $k$ -element subset  $S' \subseteq S$  the  $(S, k)$ -*private neighborhood*  $\text{pn}_k(S', S)$  of  $S'$  is the set of all vertices  $v \in V$  such that  $N(v) \cap S = S'$ . Further, the *open  $k$ -boundary*  $\text{OB}_k(S)$  of  $S$  is the set of all vertices  $v$  in  $G$  such that  $v \in \text{pn}_k(S', S)$  for some  $k$ -element subset  $S' \subseteq S$  [4]. Obviously,  $\text{OB}_k(S) = \bigcup_{S'} \text{pn}_k(S', S)$ , where  $S'$  is a  $k$ -element subset of  $S$ .

---

MSC(2010): Primary: 05C69

Keywords:  $k$ -tuple total dominating set,  $k$ -tuple total domination number,  $m$ -Mycieleskian graph.

Received: 01 November 2011, Accepted: 25 November 2011.

As we will see, the generalized Mycielskian graphs, which are also called *cones over graphs* [7], are natural generalization of Mycielski graphs. If  $V(G) = V^0 = \{v_1^0, v_2^0, \dots, v_n^0\}$  and  $E(G) = E_0$ , then for any integer  $m \geq 1$  the  $m$ -Mycielskian  $\mu_m(G)$  of  $G$  is the graph with vertex set  $V^0 \cup V^1 \cup V^2 \cup \dots \cup V^m \cup \{u\}$ , where  $V^i = \{v_j^i \mid v_j^0 \in V^0\}$  is the  $i$ -th distinct copy of  $V^0$ , for  $i = 1, 2, \dots, m$ , and edge set  $E_0 \cup \left( \bigcup_{i=0}^{m-1} \{v_j^i v_{j'}^{i+1} \mid v_j^0 v_{j'}^0 \in E_0\} \right) \cup \{v_j^m u \mid v_j^m \in V^m\}$ . The 1-Mycielskian  $\mu_1(G)$  of  $G$  is the well-studied *Mycielskian* of  $G$ , and denoted simply by  $\mu(G)$  or  $M(G)$ .

For positive integer  $k$ , the  $k$ -join of a graph  $G$  to a graph  $H$  of order at least  $k$  is the graph obtained from the disjoint union of  $G$  and  $H$  by joining each vertex of  $G$  to at least  $k$  vertices of  $H$ . We denote the  $k$ -join of  $G$  to  $H$  by  $G \circ_k H$ .

Domination in graphs is now well-studied in graph theory and the literature on this subject has been surveyed and detailed in the two books by Haynes, Hedetniemi, and Slater [2, 3].

In [4], Henning and Kazemi introduced the  $k$ -tuple total domination number of a graph. Let  $k$  be a positive integer. A subset  $S$  of  $V$  is a  $k$ -tuple total dominating set of  $G$ , abbreviated kTDS, if for every vertex  $v \in V$ ,  $|N(v) \cap S| \geq k$ , that is,  $S$  is a kTDS of  $G$  if every vertex of  $V$  has at least  $k$  neighbors in  $S$ . The  $k$ -tuple total domination number  $\gamma_{\times k, t}(G)$  of  $G$  is the minimum cardinality of a kTDS of  $G$ . We remark that a 1-tuple total domination is the well-studied *total domination number*. Thus,  $\gamma_t(G) = \gamma_{\times 1, t}(G)$ . For a graph to have a  $k$ -tuple total dominating set, its minimum degree is at least  $k$ . Since every  $(k+1)$ -tuple total dominating set is also a  $k$ -tuple total dominating set, we note that  $\gamma_{\times k, t}(G) \leq \gamma_{\times (k+1), t}(G)$  for all graphs with minimum degree at least  $k+1$ . A kTDS in a graph  $G$  is a *minimal* kTDS if no proper subset of it is a kTDS in  $G$ . A kTDS of cardinality  $\gamma_{\times k, t}(G)$  is called a  $\gamma_{\times k, t}(G)$ -set. A 2-tuple total dominating set is called a *double total dominating set*, abbreviated DTDS, and the 2-tuple total domination number is called the *double total domination number*. The redundancy involved in  $k$ -tuple total domination makes it useful in many applications. The references [5, 6] give more information about the  $k$ -tuple total domination number of a graph.

In this paper, we study the  $k$ -tuple total domination number of the  $m$ -Mycielskian graph of a graph  $G$ . We prove that for every positive integers  $m$  and  $k$  and every graph  $G$  with  $\delta(G) \geq k$ , if  $m-1 \cong r \pmod{4}$ , where  $0 \leq r \leq 3$ , and  $r' \cong r+1 \pmod{2}$ , then

$$\gamma_{\times k, t}(G) + 1 \leq \gamma_{\times k, t}(\mu_m(G)) \leq \begin{cases} (1 + 2\lceil(m-1)/4\rceil)\gamma_{\times k, t}(G) + kr' & \text{if } r = 0, 3, \\ 2\lceil(m-1)/4\rceil\gamma_{\times k, t}(G) + kr' & \text{otherwise.} \end{cases}$$

Hence  $\gamma_{\times k, t}(G) + 1 \leq \gamma_{\times k, t}(M(G)) \leq \gamma_{\times k, t}(G) + k$ . We also prove that the bounds  $\gamma_{\times k, t}(G) + 1$  and  $\gamma_{\times k, t}(G) + k$  are sharp and characterize graphs with  $\gamma_{\times k, t}(M(G)) = \gamma_{\times k, t}(G) + 1$ .

Through of this paper,  $k$  is a positive integer. The next results are useful for our investigations.

**Proposition 1.1.** (Henning, Kazemi [4] 2010) *Let  $G$  be a graph of order  $n$  with  $\delta(G) \geq k \geq 1$ , and let  $S$  be a kTDS in  $G$ . Then*

1.  $k+1 \leq \gamma_{\times k, t}(G) \leq n$ ,
2. for every spanning subgraph  $H$  of  $G$ ,  $\gamma_{\times k, t}(G) \leq \gamma_{\times k, t}(H)$ ,
3. for every vertex  $v$  of degree  $k$ ,  $N_G(v) \subseteq S$ .

**Proposition 1.2.** (Henning, Kazemi [4] 2010) *Let  $G$  be a graph of order  $n$  with  $\delta(G) \geq k \geq 1$ , and let  $S$  be a  $k$ TDS in  $G$ . Then  $S$  is a minimal  $k$ TDS of  $G$  if and only if for each vertex  $v \in S$ , there exists a  $k$ -element subset  $S_v \subseteq S$  such that  $v \in S_v$  and  $|\text{pn}_k(S_v, S)| \geq 1$ .*

**Proposition 1.3.** (Henning, Kazemi [4] 2010) *Let  $G$  be a graph with  $\delta(G) \geq k \geq 1$ . Then,  $\gamma_{\times k,t}(G) = k + 1$  if and only if  $G = K_{k+1}$  or  $G = F \circ_k K_{k+1}$  for some graph  $F$ .*

**Proposition 1.4.** (Henning, Kazemi [5] 2010) *Let  $G$  be a graph of order  $n$  with  $\delta(G) \geq k \geq 1$ . Then  $\gamma_{\times k,t}(G) \geq \lceil kn/\Delta(G) \rceil$ .*

## 2. $m$ -mycieleskian graphs

In the next theorem we give a lower bound and an upper bound on the  $k$ -tuple domination number of the  $m$ -Mycieleskian graph  $\mu_m(G)$  in terms  $k$  and the  $k$ -tuple domination number of  $G$ . First we state the following lemma which has an easy proof that is left to the reader.

**Lemma 2.1.** *Let  $G$  be a graph with  $\delta(G) \geq k \geq 1$ . Let  $V(\mu_m(G)) = V^0 \cup V^1 \cup V^2 \cup \dots \cup V^m \cup \{u\}$ . If  $\gamma_{\times k,t}(\mu_m(G)) = \gamma_{\times k,t}(G)$ , then for every  $\gamma_{\times k,t}(\mu_m(G))$ -set  $S$ ,  $u \notin S$ , and so  $m = 1$ .*

**Theorem 2.2.** *Let  $m$  and  $k$  be two positive integers, and let  $G$  be a graph with  $\delta(G) \geq k \geq 1$ . Then*

$$\gamma_{\times k,t}(G) + 1 \leq \gamma_{\times k,t}(\mu_m(G)) \leq \begin{cases} (1 + 2\lceil(m-1)/4\rceil)\gamma_{\times k,t}(G) & \text{if } m \cong 0 \pmod{4}, \\ (1 + 2\lceil(m-1)/4\rceil)\gamma_{\times k,t}(G) + k & \text{if } m \cong 1 \pmod{4}, \\ 2\lceil(m-1)/4\rceil\gamma_{\times k,t}(G) & \text{if } m \cong 2 \pmod{4}, \\ 2\lceil(m-1)/4\rceil\gamma_{\times k,t}(G) + k & \text{if } m \cong 3 \pmod{4}. \end{cases}$$

*Proof.* Let  $V(\mu_m(G)) = V^0 \cup V^1 \cup V^2 \cup \dots \cup V^m \cup \{u\}$ . Since  $G$  is an induced subgraph of  $\mu_m(G)$ ,  $\gamma_{\times k,t}(G) \leq \gamma_{\times k,t}(\mu_m(G))$ . If  $\gamma_{\times k,t}(\mu_m(G)) = \gamma_{\times k,t}(G)$ , then Lemma 2.1 implies  $m = 1$  and for every  $\gamma_{\times k,t}(\mu_m(G))$ -set  $S$ ,  $u \notin S$ . Since every vertex of  $V^1$  is adjacent to at least  $k$  vertices of  $S \cap V^0$ , we conclude that every vertex of  $V^0$  is adjacent to at least  $k$  vertices of  $S \cap V^0$ . Hence

$$\gamma_{\times k,t}(G) \leq |S \cap V^0| = |S| - |S \cap V^1| \leq \gamma_{\times k,t}(G) - k < \gamma_{\times k,t}(G),$$

a contradiction. Therefore  $\gamma_{\times k,t}(\mu_m(G)) \geq \gamma_{\times k,t}(G) + 1$ .

Now we prove the other inequality. For an arbitrary  $\gamma_{\times k,t}(G)$ -set  $S$ , let  $S^i = \{v^i \mid v \in S\} \subseteq V^i$  be the  $i$ -th distinct copy of  $S$  when  $0 \leq i \leq m$ . Let also  $S_k$  be an arbitrary subset of  $V^m$  of cardinality  $k$ . We continue our proof in the following four cases.

**Case 0.**  $m \cong 0 \pmod{4}$ .

The set  $S' = S^0 \cup (\bigcup_{t=1}^{\lfloor(m-1)/4\rfloor} (S^{4t-1} \cup S^{4t})) \cup (S^{m-1} \cup S^m)$  is a  $k$ TDS of  $\mu_m(G)$  of cardinality

$$(3 + 2\lfloor(m-1)/4\rfloor)\gamma_{\times k,t}(G) = (1 + 2\lceil(m-1)/4\rceil)\gamma_{\times k,t}(G).$$

**Case 1.**  $m \cong 1 \pmod{4}$ .

The set  $S' = S^0 \cup (\bigcup_{t=1}^{\lfloor(m-1)/4\rfloor} (S^{4t-1} \cup S^{4t})) \cup S_k$  is a  $k$ TDS of  $\mu_m(G)$  of cardinality

$$(1 + 2\lfloor(m-1)/4\rfloor)\gamma_{\times k,t}(G) + k = (1 + 2\lceil(m-1)/4\rceil)\gamma_{\times k,t}(G) + k.$$

**Case 2.**  $m \cong 2 \pmod{4}$ .

The set  $S' = (\bigcup_{t=1}^{\lfloor (m-1)/4 \rfloor} (S^{4t-3} \cup S^{4t-2})) \cup (S^{m-1} \cup S^m)$  is a kTDS of  $\mu_m(G)$  of cardinality

$$(2 + 2\lfloor (m-1)/4 \rfloor)\gamma_{\times k,t}(G) = 2\lceil (m-1)/4 \rceil\gamma_{\times k,t}(G).$$

**Case 3.**  $m \cong 3 \pmod{4}$ .

The set  $S' = (\bigcup_{t=1}^{\lceil (m-1)/4 \rceil} (S^{4t-3} \cup S^{4t-2})) \cup S_k$  is a kTDS of  $\mu_m(G)$  of cardinality

$$2\lceil (m-1)/4 \rceil\gamma_{\times k,t}(G) + k.$$

Therefore we have proved

$$\gamma_{\times k,t}(\mu_m(G)) \leq |S'| = \begin{cases} (1 + 2\lceil (m-1)/4 \rceil)\gamma_{\times k,t}(G) & \text{if } m \cong 0 \pmod{4}, \\ (1 + 2\lceil (m-1)/4 \rceil)\gamma_{\times k,t}(G) + k & \text{if } m \cong 1 \pmod{4}, \\ 2\lceil (m-1)/4 \rceil\gamma_{\times k,t}(G) & \text{if } m \cong 2 \pmod{4}, \\ 2\lceil (m-1)/4 \rceil\gamma_{\times k,t}(G) + k & \text{if } m \cong 3 \pmod{4}, \end{cases}$$

and this completes our proof.  $\square$

**Corollary 2.3.** *Let  $m$  be a positive integer, and let  $G$  be a graph without isolated vertex. Then*

$$\gamma_t(G) + 1 \leq \gamma_t(\mu_m(G)) \leq \begin{cases} (1 + 2\lceil (m-1)/4 \rceil)\gamma_t(G) & \text{if } m \cong 0 \pmod{4}, \\ (1 + 2\lceil (m-1)/4 \rceil)\gamma_t(G) + 1 & \text{if } m \cong 1 \pmod{4}, \\ 2\lceil (m-1)/4 \rceil\gamma_t(G) & \text{if } m \cong 2 \pmod{4}, \\ 2\lceil (m-1)/4 \rceil\gamma_t(G) + 1 & \text{if } m \cong 3 \pmod{4}. \end{cases}$$

### 3. Mycielekian graphs

Theorem 2.2 implies the next two theorems when  $m = 1$ .

**Theorem 3.1.** *If  $G$  is a graph with no isolated vertices, then  $\gamma_t(M(G)) = \gamma_t(G) + 1$ .*

**Theorem 3.2.** *If  $G$  is a graph with  $\delta(G) \geq k \geq 2$ , then  $\gamma_{\times k,t}(G) + 1 \leq \gamma_{\times k,t}(M(G)) \leq \gamma_{\times k,t}(G) + k$ .*

In the next theorem we give other lower bound for  $\gamma_{\times k,t}(M(G))$ .

**Theorem 3.3.** *If  $G$  is a graph with  $\delta(G) \geq k \geq 2$ , then*

$$\gamma_{\times k,t}(M(G)) \geq \min\{\gamma_{\times k,t}(G) + k, \gamma_{\times(k-1),t}(G) + k + 1\}.$$

*Proof.* Let  $V(M(G)) = V^0 \cup V^1 \cup \{u\}$  and let  $S$  be an arbitrary kTDS of  $M(G)$ . Then  $|S \cap V^1| \geq k$ . If  $u \in S$ , then  $|S \cap V^0| \geq \gamma_{\times(k-1),t}(G)$ . Since  $S \cap V^0$  must be a  $(k-1)$ TDS of  $V^1$ . If  $u \notin S$ , then  $|S \cap V^0| \geq \gamma_{\times k,t}(G)$ . Since  $S \cap V^0$  must be a kTDS of  $V^1$ . Therefore

$$\gamma_{\times k,t}(M(G)) \geq \min\{\gamma_{\times k,t}(G) + k, \gamma_{\times(k-1),t}(G) + k + 1\}.$$

$\square$

As an immediately result of Theorems 3.2 and 3.3 we have the following two corollaries.

**Corollary 3.4.** *Let  $G$  be a graph with  $\delta(G) \geq k \geq 2$ . If  $\gamma_{\times k,t}(G) = \gamma_{\times(k-1),t}(G) + 1$ , then*

$$\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + k.$$

**Corollary 3.5.** *If  $n \geq k + 1 \geq 3$ , then  $\gamma_{\times k,t}(M(K_n)) = \gamma_{\times k,t}(K_n) + k = 2k + 1$ .*

Also the following two results show that the upper bound  $\gamma_{\times k,t}(G) + k$  in Theorem 3.2 is sharp for some of the complete multipartite graphs, the complete graph  $K_{k+1}$  and the  $k$ -join  $F \circ_k K_{k+1}$ , for every graph  $F$ .

**Proposition 3.6.** *Let  $G = K_{n_1, \dots, n_p}$  be a complete  $p$ -partite graph. If  $p \geq k + 1 \geq 3$ , then*

$$\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + k.$$

*Proof.* We suppose that  $V(G)$  has partition  $X_1 \cup X_2 \cup \dots \cup X_p$  such that  $|X_j| = n_j$  for  $j = 1, 2, \dots, p$ . Let  $V(M(G)) = V^0 \cup V^1 \cup \{u\}$ , where  $V^i = X_1^i \cup X_2^i \cup \dots \cup X_p^i$  and  $X_j^i = \{v_j^i \mid v_j \in X_j\}$ , for  $i = 0, 1$  and  $j = 1, 2, \dots, p$ . Let also  $S$  be an arbitrary kTDS of  $M(G)$ . Obviously  $|S \cap V^1| \geq k$ , and, without loss of generality, we may assume that  $|S \cap V^1| = k$ . Let  $S \cap V^1$  be a set which contains only one vertex of every  $X_j^1$ , for  $1 \leq j \leq k$ . Thus each vertex of  $V^0 - \{v_i^0 \mid v_i^1 \in S \cap V^1\}$  is adjacent to all vertices in  $S \cap V^1$ . Since each vertex of  $S \cap V^1$  must be adjacent to at least  $k$  vertices of  $S$ , we have  $|S \cap (V^0 \cup \{u\})| \geq k$ . The assumptions  $k \geq 2$  and  $|S \cap V^0| \geq k - 1$  imply  $S \cap V^0 \neq \emptyset$ . We see that there exists a unique index  $1 \leq j \leq k$  such that each vertex of  $X_j^1$  is adjacent to  $k - 1$  vertices of  $S \cap (V^0 \cup \{u\})$ . Hence  $|S \cap (V^0 \cup \{u\})| \geq k + 1$ , and so

$$\begin{aligned} \gamma_{\times k,t}(M(G)) &= \min \{|S| : S \text{ is a kTDS of } M(G)\} \\ &\geq 2k + 1 \\ &= \gamma_{\times k,t}(G) + k. \end{aligned}$$

Now Theorem 3.2 implies  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + k$ . □

**Theorem 3.7.** *Let  $G$  be a graph of order  $n \geq k + 1$  with  $\delta(G) \geq k \geq 2$ . If  $G$  is the  $k$ -join  $F \circ_k K_{k+1}$ , for some graph  $F$ , then*

$$\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + k.$$

*Proof.* Let  $G$  be the  $k$ -join  $F \circ_k K_{k+1}$ , for some graph  $F$ . Then Proposition 1.3 implies  $\gamma_{\times k,t}(G) = k + 1$ . Since  $G$  is a spanning subgraph of  $K_n$ , and hence  $M(G)$  is a spanning subgraph of  $M(K_n)$ , we have

$$\begin{aligned} \gamma_{\times k,t}(M(G)) &\geq \gamma_{\times k,t}(M(K_n)) \\ &= \gamma_{\times k,t}(K_n) + k \\ &= 2k + 1 \\ &= \gamma_{\times k,t}(G) + k. \end{aligned}$$

Now Theorem 3.2 implies  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + k = 2k + 1$ . □

Theorem 3.1 shows that  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + 1$ , where  $G$  is a graph with no isolated vertices and  $k = 1$ . Here, we give an equivalent condition for  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + 1$ , when  $k \geq 2$ . We recall that a kTDS  $S$  is a minimal kTDS if and only if for each vertex  $v \in S$ , there exists a  $k$ -element subset  $S_v \subseteq S$  such that  $v \in S_v$  and  $|\text{pn}_k(S_v, S)| \geq 1$ .

**Theorem 3.8.** *Let  $G$  be a graph with  $\delta(G) \geq k \geq 1$ . Then  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + 1$  if and only if  $k = 1$  or  $k \geq 2$  and  $G$  has a  $\gamma_{\times k,t}$ -set  $S$  with a  $k$ -subset  $S' \subseteq S$  such that  $S - S'$  is a  $(k - 1)$ TDS of  $G$  and for every vertex  $v$ ,  $|S_v \cap S'| \leq 1$ .*

*Proof.* Let  $V(M(G)) = V^0 \cup V^1 \cup \{u\}$ , where  $V^i = \{v_j^i \mid 1 \leq j \leq n\}$  for  $i = 0, 1$ . If  $k = 1$ , then  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + 1$ , by Theorem 3.1. Now let  $k \geq 2$ . Let  $S$  be a  $\gamma_{\times k,t}(G)$ -set which contains a  $k$ -subset  $S' \subseteq S$  with this conditions that  $S - S'$  is a  $(k - 1)$ TDS of  $G$  and for every vertex  $v$ ,  $|S_v \cap S'| \leq 1$ . Since

$$D = \{v_j^0 \mid v_j \in S - S'\} \cup \{v_j^1 \mid v_j \in S'\} \cup \{u\}$$

is a kTDS of  $M(G)$  of cardinality  $\gamma_{\times k,t}(G) + 1$ , Theorem 3.2 implies  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + 1$ .

Conversely, let  $\gamma_{\times k,t}(M(G)) = \gamma_{\times k,t}(G) + 1$  and let  $k \geq 2$ . If  $u$  belongs to a  $\gamma_{\times k,t}$ -set of  $M(G)$ , we have no thing to prove. Thus we assume that  $u$  belongs to no  $\gamma_{\times k,t}(M(G))$ -set  $D$ , and so  $|D \cap V^1| \geq k$  and  $|D \cap V^0| \geq \gamma_{\times k,t}(G)$ . Therefore

$$\gamma_{\times k,t}(G) + 1 = \gamma_{\times k,t}(M(G)) = |D| \geq \gamma_{\times k,t}(G) + k \geq \gamma_{\times k,t}(G) + 2,$$

a contradiction.  $\square$

The next proposition gives graphs that satisfy in the condition of Theorem 3.8. First, we present the definition of the *Harary graph* [8].

Given  $m \leq n$ , place  $n$  vertices  $1, 2, \dots, n$  around a circle, equally spaced. If  $m$  is even, form  $H_{m,n}$  by making each vertex adjacent to the nearest  $m/2$  vertices in each direction around the circle. If  $m$  is odd and  $n$  is even, form  $H_{m,n}$  by making each vertex adjacent to the nearest  $(m - 1)/2$  vertices in each direction and to the diametrically opposite vertex. In each case,  $H_{m,n}$  is  $m$ -regular. When  $m$  and  $n$  are both odd, index the vertices by the integers modulo  $n$ . Construct  $H_{m,n}$  from  $H_{m-1,n}$  by adding the edges  $i \leftrightarrow i + (n - 1)/2$  for  $0 \leq i \leq (n - 1)/2$ .

**Proposition 3.9.** *If  $G$  is a cycle of order at least 3 or the Harary graph  $H_{2m,\ell m+1}$ , where  $\ell \geq 3$  and  $m \geq 1$ , then*

$$\gamma_{\times 2,t}(M(G)) = \gamma_{\times 2,t}(G) + 1.$$

*Proof.* Let  $G = C_n$  be a cycle of order at least 3 with the vertex set  $V(C_n) = V^0 = \{v_j \mid 1 \leq j \leq n\}$  and the edge set  $E(C_n) = \{(v_j, v_{j+1}) \mid 1 \leq j \leq n\}$ . Let also  $V(M(G)) = V^0 \cup V^1 \cup \{u\}$ . Proposition 1.1(3) implies  $\gamma_{\times 2,t}(G) = n$ . Since  $S = (V^0 - \{v_1^0, v_n^0\}) \cup \{v_{n-1}^1, v_n^1, u\}$  is a DTDS of  $M(G)$  of cardinality  $\gamma_{\times 2,t}(C_n) + 1 = n + 1$ , Theorem 2.2 implies  $\gamma_{\times 2,t}(M(C_n)) = \gamma_{\times 2,t}(C_n) + 1 = n + 1$ .

Now let  $G = H_{2m,\ell m+1}$  and let  $V(M(G)) = V^0 \cup V^1 \cup \{u\}$ , where  $V^i = \{v_j^i \mid 1 \leq j \leq \ell m + 1\}$  for  $i = 0, 1$ ,  $V^0 = V(G)$  and  $\ell \geq 3$ . Since  $\{v_{jm+1}^0 \mid 0 \leq j \leq \lceil (\ell m + 1)/m \rceil - 1 = \ell\}$  is a DTDS of  $G$ , we have  $\gamma_{\times 2,t}(G) = \lceil (\ell m + 1)/m \rceil = \ell + 1$ , by Proposition 1.4. Since also  $S = \{v_{im+1}^0 \mid 1 \leq i \leq \ell - 1\} \cup \{v_1^1, v_{\ell m+1}^1, u\}$  is a DTDS of  $M(G)$  of cardinality  $\ell + 2 = \gamma_{\times 2,t}(G) + 1$ , Theorem 2.2 implies  $\gamma_{\times 2,t}(M(G)) = \gamma_{\times 2,t}(G) + 1$ .  $\square$

In the end of paper, the author states the following problem.

**Problem:** For integers  $k, m \geq 1$ , characterize graphs  $G$  with  $\delta(G) \geq k$  satisfy  $\gamma_{\times k,t}(\mu_m(G)) = \gamma_{\times k,t}(G) + 1$  or

$$\gamma_{\times k,t}(\mu_m(G)) = \begin{cases} (1 + 2\lceil(m-1)/4\rceil)\gamma_{\times k,t}(G) & \text{if } m \equiv 0 \pmod{4}, \\ (1 + 2\lceil(m-1)/4\rceil)\gamma_{\times k,t}(G) + k & \text{if } m \equiv 1 \pmod{4}, \\ 2\lceil(m-1)/4\rceil\gamma_{\times k,t}(G) & \text{if } m \equiv 2 \pmod{4}, \\ 2\lceil(m-1)/4\rceil\gamma_{\times k,t}(G) + k & \text{if } m \equiv 3 \pmod{4}. \end{cases}$$

REFERENCES

[1] F. Harary, T. W. Haynes, Double domination in graphs, *Ars Combin.* **55** (2000) 201–213.  
 [2] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, *Fundamentals of Domination in Graphs*, Marcel Dekker, New York, 1998.  
 [3] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, *Domination in Graphs: Advanced Topics*, Marcel Dekker, New York, 1998.  
 [4] M. A. Henning, A. P. Kazemi,  $k$ -tuple total domination in graphs, *Discrete Applied Mathematics* **158** (2010) 1006–1011.  
 [5] M. A. Henning, A. P. Kazemi,  $k$ -tuple total domination in cross product graphs, *Journal of Combinatorial Optimization* DOI 10.1007/s10878-011-9389-z.  
 [6] A. P. Kazemi,  $k$ -tuple total domination in complementary prisms, To appear in *ISRN Discrete Mathematics*.  
 [7] C. Tardif, Fractional chromatic numbers of cones over graphs, *J. Graph Theory* **38** (2001) 87–94.  
 [8] D. B. West, *Introduction to Graph Theory*, (2nd edition), Prentice Hall, USA, 2001.

**Adel P. Kazemi**

Department of Mathematics University of Mohaghegh Ardabili, P. O. Box 5619911367 Ardabil, Iran

Email: a.p.kazemi@uma.ac.ir

Email: adelpkazemi@yahoo.com