

**Transactions on Combinatorics** ISSN (print): 2251-8657, ISSN (on-line): 2251-8665 Vol. 4 No. 1 (2015), pp. 49-56. © 2015 University of Isfahan



# k-ODD MEAN LABELING OF PRISM

B. GAYATHRI\* AND K. AMUTHAVALLI

Communicated by Jamshid Moori

ABSTRACT. A (p,q) graph G is said to have a k-odd mean labeling  $(k \ge 1)$  if there exists an injection  $f: V \to \{0, 1, 2, \dots, 2k + 2q - 3\}$  such that the induced map  $f^*$  defined on E by  $f^*(uv) = \left\lceil \frac{f(u) + f(v)}{2} \right\rceil$  is a bijection from E to  $\{2k - 1, 2k + 1, 2k + 3, \dots, 2k + 2q - 3\}$ . A graph that admits k-odd mean labeling is called k-odd mean graph. In this paper, we investigate k-odd mean labeling of prism  $C_m \times P_n$ .

## 1. Introduction

All graphs in this paper are finite, simple and undirected. Terms not defined here are used in the sense of Harary [11]. A graph labeling is an assignment of integers to the vertices or edges or both subject to certain conditions. If the domain of the mapping is the set of vertices (or edges) then the labeling is called a vertex labeling (or an edge labeling). Labeled graphs serve as useful models for a broad range of applications such as X-ray crystallography, radar, coding theory, astronomy, circuit design and communication network addressing. Graph labeling was first introduced in the late 1960s. Many studies in graph labeling refer to Rosas research in 1967 [13]. Mean labeling of graphs was discussed in [14] and the concept of odd mean labeling was introduced in [12]. In [6], we introduced k-odd mean labeling to (k, d)-odd mean labeling. In this paper, we investigate k-odd mean labeling of Prism  $(C_m \times P_n)$ .

### 2. Definitions

**Definition 2.1** (k-odd mean labeling). A (p,q) graph G is said to have a k-odd mean labeling  $(k \ge 1)$  if there exists a injection  $f: V \to \{0, 1, 2, \dots, 2k + 2q - 3\}$  such that the induced map  $f^*$  defined on E by  $f^*(uv) = \left\lceil \frac{f(u)+f(v)}{2} \right\rceil$  is an bijection from E to  $\{2k-1, 2k+1, 2k+3, \dots, 2k+2q-3\}$ .

MSC(2010): 05C78.

Keywords: k-odd mean labeling, k-odd mean graph.

Received: 16 September 2013, Accepted: 12 May 2014.

<sup>\*</sup>Corresponding author.

A graph that admits a k-odd mean labeling is called a k-odd mean graph.

**Definition 2.2** ((k, d)-odd mean labeling). A (p,q) graph G is said to have a (k,d)-odd mean labeling ( $k \ge 1$  and  $d \ge 1$ ) if there exists a injection  $f: V \to \{0, 1, 2, \dots, 2k - 1 + 2(q - 1)d\}$  such that the induced map  $f^*$  defined on E by  $f^*(uv) = \left\lceil \frac{f(u)+f(v)}{2} \right\rceil$  is an bijection from E to  $\{2k-1, 2k-1+2d, 2k-1+4d, \dots, 2k-1+2(q-1)d\}$ .

A graph that admits a (k, d)-odd mean labeling is called a (k, d)-odd mean graph.

**Remark 2.3.** 1-odd mean labeling is an odd mean labeling.

**Remark 2.4.** The graphs  $C_3$  and  $C_6$  are not k-odd mean graphs.

**Definition 2.5.** The Cartesian product  $G_1 \times G_2$  of two graphs  $G_1$  and  $G_2$  is the simple graph with  $V_1 \times V_2$  as its vertex set and two vertices  $(u_1, v_1)$  and  $(u_2, v_2)$  are adjacent in  $G_1 \times G_2$  iff either  $u_1 = u_2$  and  $v_1$  is adjacent to  $v_2$  in  $G_2$ , or  $u_1$  is adjacent to  $u_2$  in  $G_1$  and  $v_1 = v_2$ .

**Remark 2.6.** For brevity, we use k-OML for k-odd mean labeling.

#### 3. Main result

**Theorem 3.1.**  $C_m \times P_n$   $(m \ge 4, n \ge 2)$  is a k-odd mean graph for any k and  $m \ne 6$ .

Proof.

$$V(C_m \times P_n) = \{v_{ij}, 1 \le i \le n \text{ and } 1 \le j \le m\} \text{ and}$$
$$E(C_m \times P_n) = \{e_{ij}, 1 \le i \le n \text{ and } 1 \le j \le m\}$$
$$\cup \{e'_{ij}, 1 \le i \le n-1 \text{ and } 1 \le j \le m\} \text{ (see Fig. 1)}.$$



FIGURE 1. Ordinary labeling of  $C_m \times P_n$ 

First we label the vertices of  $C_m \times P_n$  as follows:

Define  $f: V(C_m \times P_n) \rightarrow \{0, 1, 2, \dots, 2k + 2q - 3\}$  by **Case (i)**  $m \equiv 0 \pmod{4}$ . For  $1 \leq i \leq n$  and i is odd, for  $1 \leq j \leq \frac{m}{2}$ ,

$$f(v_{ij}) = \begin{cases} 2k + 4m(i-1) + 4j - 6, & \text{if } j \text{ is odd,} \\ 2k + 4m(i-1) + 4j - 8, & \text{if } j \text{ is even} \end{cases}$$

for  $\frac{m+2}{2} \le j \le m$ ,

$$f(v_{ij}) = \begin{cases} 2k + 4(mi - j) + 1, & \text{if } j \text{ is odd,} \\ 2k + 4(mi - j + 1), & \text{if } j \text{ is even} \end{cases}$$

For  $1 \le i \le n$  and i is even, for  $1 \le j \le \frac{m-2}{2}$ ,

$$f(v_{ij}) = \begin{cases} 2k + 4m(i-1) + 4(j-1), & \text{if } j \text{ is odd,} \\ 2k + 4m(i-1) + 4(j-1) - 2, & \text{if } j \text{ is even} \end{cases}$$

for  $\frac{m}{2} \leq j \leq m$ ,

$$f(v_{ij}) = \begin{cases} 2k + 4(mi - j), & \text{if } j \text{ is odd,} \\ 2k + 4(mi - j) - 3, & \text{if } j \text{ is even.} \end{cases}$$

Then the induced edge labels are:

For  $1 \leq i \leq n$  and i is odd,

$$f^*(e_{ij}) = 2k + 4m(i-1) + 4j - 5, \ 1 \le j \le \frac{m}{2},$$
  
$$f^*\left(e_{i\frac{m+2}{2}}\right) = 2k + 2m + 4m(i-1) - 3,$$
  
$$f^*(e_{ij}) = 2k + 4mi - 4j + 1, \ \frac{m+4}{2} \le j \le m.$$

For  $1 \leq i \leq n$  and i is even,

$$f^*(e_{ij}) = \begin{cases} 2k + 4m(i-1) + 4(j-1) + 3, & 1 \le j \le \frac{m-2}{2}, \\ 2k + 4mi - 4j - 3, & \frac{m}{2} \le j \le m-1. \end{cases}$$
$$f^*(e_{im}) = 2k + 4m(i-1) - 1.$$

For  $1 \leq i \leq n-1$ ,

$$f^*(e'_{ij}) = \begin{cases} 2k + 2m(2i-1) + 4j - 5, & 1 \le j \le \frac{m}{2}, \\ 2k + 2m(2i-1) + 4j - 3, & \frac{m+2}{2} \le j \le m-1. \end{cases}$$

3-OML of  $C_8 \times P_3$  is shown in Fig. 2.



FIGURE 2. 3-OML of  $C_8 \times P_3$ .

Case (ii)  $m \equiv 1, 3 \pmod{4}$ .

The vertex labels are:

For  $1 \le i \le n$  and i is odd, for  $1 \le j \le \frac{m-3}{2}$ ,

$$f(v_{ij}) = 2k + 4m(i-1) + 2(j-2),$$
  
$$f\left(v_{i\frac{m-1}{2}}\right) = 2k + m(4i-3) - 6,$$
  
$$f\left(v_{i\frac{m+1}{2}}\right) = 2k + m(4i-3) - 2$$

for  $\frac{m+3}{2} \le j \le m-1$ ,

$$f(v_{ij}) = 2k + 4m(i-1) + 2(j-1),$$
  
$$f(v_{im}) = 2k + 2m(2i-1) - 3.$$

For  $1 \le i \le n$  and i is even, for  $1 \le j \le \frac{m-5}{2}$ ,

$$f(v_{ij}) = 2k + 4m(i-1) + 2(j-1),$$
  
$$f\left(v_{i\frac{m-1}{2}}\right) = 2k + m(4i-3) - 2$$

for  $\frac{m+1}{2} \le j \le m-2$ ,

$$f(v_{ij}) = 2k + 4m(i-1) + 2j,$$
  
$$f(v_{im-1}) = 2k + 4m(i-1) + 2n - 3,$$
  
$$f(v_{im}) = 2k + 4m(i-1) - 2.$$

Then the induced edge labels are: For  $1 \le i \le n$  and i is odd,

$$f^*(e_{ij}) = \begin{cases} 2k + 4m(i-1) + 2j - 3, 1 \le j \le \frac{m-1}{2}, \\ 2k + 4m(i-1) + 2j - 1, \frac{m+1}{2} \le j \le m-1 \end{cases}$$
$$f^*(e_{im}) = 2k + m(4i - 3) - 2.$$

For  $1 \le i \le n$  and i is even, for  $1 \le j \le \frac{m-3}{2}$ ,

$$f^*(e_{ij}) = 2k + 4m(i-1) + 2j - 1$$

for  $\frac{m-1}{2} \le j \le m-2$ ,

$$f^*(e_{ij}) = 2k + 4m(i-1) + 2j + 1,$$
  
$$f^*(e_{im-1}) = 2k + m(4i-3) - 2,$$
  
$$f^*(e_{im}) = 2k + 4m(i-1) - 1.$$

For  $1 \le i \le n-1$  and  $1 \le j \le \frac{m-1}{2}$ ,

$$f^*(e'_{ij}) = 2k + 2m(2i - 1) + 2j - 3$$

for  $\frac{m+1}{2} \le j \le m-1$ ,

$$f^*(e'_{ij}) = 2k + 2m(2i - 1) + 2j - 1,$$
  
$$f^*(e'_{im}) = 2k + m(4i - 1) - 2.$$

4-OML of  $C_5 \times P_5$  is shown Fig. 3.



FIGURE 3. 4-OML of  $C_5 \times P_5$ .

5-OML of  $C_{11} \times P_3$  is shown in Fig. 4.



FIGURE 4. 5-OML of  $C_{11} \times P_3$ .

Case (iii)  $m \equiv 2 \pmod{4}$  and m > 6. The vertex labels are: For  $1 \le i \le n$  and i is odd, for  $1 \le j \le \frac{m-4}{2}$ ,

$$f(v_{ij}) = 2k + 4m(i-1) + 2(j-2),$$
  
$$f\left(v_{i\frac{m-2}{2}}\right) = 2k + m(4i-3) - 7,$$
  
$$f\left(v_{i\frac{m}{2}}\right) = 2k + m(4i-3) - 3$$

for  $\frac{m+2}{2} \le j \le m-3$ ,

$$f(v_{ij}) = 2k + 4m(i-1) + 2(j-1),$$
  

$$f(v_{im-2}) = 2k + 2m(2i-1) - 7,$$
  

$$f(v_{im-1}) = 2k + 2n(2i-1) - 3,$$
  

$$f(v_{im}) = 2k + 2m(2i-1) - 4.$$

For  $1 \le i \le n$  and i is even, for  $1 \le j \le \frac{m-6}{2}$ ,

$$\begin{split} f(v_{ij}) &= 2k + 4m(i-1) + 2(j-1), \\ f\left(v_{i\frac{m-4}{2}}\right) &= 2k + m(4i-3) - 7, \\ f\left(v_{i\frac{m-2}{2}}\right) &= 2k + m(4i-3) - 3 \end{split}$$

Trans. Comb. 4 no. 1 (2015) 49-56

for  $\frac{m}{2} \le j \le m - 4$ ,

$$f(v_{ij}) = 2k + 4m(i-1) + 2j,$$
  

$$f(v_{im-3}) = 2k + 2m(2i-1) - 7,$$
  

$$f(v_{im-2}) = 2k + 2m(2i-1) - 3,$$
  

$$f(v_{im-1}) = 2k + 2m(2i-1) - 4,$$
  

$$f(v_{im}) = 2k + 4m(i-1) - 2.$$

Then the induced edge labels are:

For  $1 \leq i \leq n$  and i is odd,

$$f^*(e_{ij}) = \begin{cases} 2k + 4m(i-1) + 2j - 3, 1 \le j \le \frac{m-2}{2}, \\ 2k + 4m(i-1) + 2j - 1, \frac{m}{2} \le j \le m-1, \end{cases}$$
$$f^*(e_{im}) = 2k + m(4i - 3) - 3.$$

For  $1 \le i \le n$  and i is even, for  $1 \le j \le \frac{m-4}{2}$ ,

for 
$$\frac{m-2}{2} \le j \le m-2$$
,  
 $f^*(e_{ij}) = 2k + 4m(i-1) + 2j - 1$   
 $f^*(e_{ij}) = 2k + 4m(i-1) + 2j + 1$ ,  
 $f^*(e_{im-1}) = 2k + m(4i-3) - 3$ ,  
 $f^*(e_{im}) = 2k + 4m(i-1) - 1$ .

For  $1 \le i \le n-1$  and  $1 \le j \le \frac{m-2}{2}$ ,

For 
$$\frac{m}{2} \le j \le m-1$$
,  
 $f^*(e'_{ij}) = 2k + 2m(2i-1) + 2j - 3.$   
 $f^*(e'_{ij}) = 2k + 2m(2i-1) + 2j - 1,$   
 $f^*(e'_{im}) = 2k + m(4i-1) - 3.$ 

2-OML of  $C_{14} \times P_3$  is shown in



FIGURE 5. 2-OML of  $C_{14} \times P_3$ .

Therefore,  $f^*(E(C_m \times P_n) = \{2k - 1, 2k + 1, 2k + 3, \dots, 2k + 2q - 3\}.$ 

So, f is a k-odd mean labeling and hence,  $C_m \times P_n$  is k-odd mean graph for any k when  $m \ge 4$ ,  $n \ge 2$  and  $m \ne 6$ .

### References

- K. Amuthavalli, Graph labeling and its applications-Some generalization of odd mean labeling, Ph. D. Thesis, Mother Teresa Womens University, 2010.
- [2] G. S. Bloom and S. W. Golomb, Numbered complete graphs, unusual rulers and assorted applications, Theory and Applications of Graphs, Lecture notes in Mathematics, 642, Springer, Berlin, 1978 53–65.
- [3] G. S. Bloom and S. W. Colomb, Applications of Numbered undirected graphs, Proceedings of IEEE, 65 (1977) 562–570.
- [4] G. S. Bloom and D. F. Hsu, On graceful digraphs and a problem in network addressing, Congr. Numer., 35 (1982) 91–103.
- [5] A. J. Gallian, A dynamic survey of graph labeling, *Electron. J. Combin.*, Dynamic Survey 6, 5 (1998) pp. 43.
- [6] B. Gayathri and K. Amuthavalli, k-odd mean labeling of graphs, Proceedings of the International Conference on Mathematics and Computer Science-2, Sitech Publications, 2007 112–115.
- [7] B. Gayathri and K. Amuthavalli, (k, d)-odd mean labeling of graphs, National Symposium on Mathematical Methods and Applications, I. I. T., Chennai, 2006.
- [8] B. Gayathri and K. Amuthavalli, k-odd mean labeling of Crown graphs, Int. J. Math. Comput. Sci., 2 no. 3 (2007) 253–259.
- [9] B. Gayathri and K. Amuthavalli, (k, d)-odd mean labeling of some graphs, Bull. Pure Appl. Sci. Sect. E Math. Stat., 26 no. 2 (2007) 263–267.
- [10] B. Gayathri and K. Amuthavalli, k-odd mean labeling of  $\langle K_{1,n}, K_{1,m} \rangle$ , Acta Cienc. Indica Math., **34** no. 2 (2008) 827–834.
- [11] F. Harary, Graph Theory, Addison Wesley, Mass Reading, 1972.
- [12] K. Manickam and M. Marudai, Odd mean labelings of graphs, Bull. Pure Appl. Sci. Sect. E Math. Stat., 25 no. 1 (2006) 149–153.
- [13] A. Rosa, On certain valuations of the vertices of a graph, Theory of Graphs Internat. Sympos., Rome, 1966), Gordon and Breach, New York; Dunod, Paris, 1967 349–355.
- [14] S. Somasundaram and R. Ponraj, Mean Labeling of graphs, Nat. Acad. Sci. Lett., 26 no. 7-8 (2003) 210-213.

#### B. Gayathri

Department of Mathematics, Periyar E. V. R. College(Autonomous), Tiruchirappalli-620 020 Email: maduraigayathri@gmail.com

#### K. Amuthavalli

Department of Mathematics, Roever Engineering College, Perambalur-621 212 Email: thrcka@gmail.com