



www.combinatorics.ir

Transactions on Combinatorics

ISSN (print): 2251-8657, ISSN (on-line): 2251-8665

Vol. 4 No. 3 (2015), pp. 25-35.

© 2015 University of Isfahan



www.ui.ac.ir

ON LAPLACIAN-ENERGY-LIKE INVARIANT AND INCIDENCE ENERGY

S. PIRZADA* AND HILAL A. GANIE

Communicated by Ivan Gutman

ABSTRACT. For a simple connected graph G with n -vertices having Laplacian eigenvalues $\mu_1, \mu_2, \dots, \mu_{n-1}, \mu_n = 0$, and signless Laplacian eigenvalues q_1, q_2, \dots, q_n , the Laplacian-energy-like invariant (LEL) and the incidence energy (IE) of a graph G are respectively defined as $LEL(G) = \sum_{i=1}^{n-1} \sqrt{\mu_i}$ and $IE(G) = \sum_{i=1}^n \sqrt{q_i}$. In this paper, we obtain some sharp lower and upper bounds for the Laplacian-energy-like invariant and incidence energy of a graph.

1. Introduction

Let G be a finite, undirected, simple graph with n vertices and m edges having vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$. The adjacency matrix $A = (a_{ij})$ of G is a $(0, 1)$ -square matrix of order n whose (i, j) -entry is equal to 1 if v_i is adjacent to v_j and equal to 0, otherwise. Let $D(G) = \text{diag}(d_1, d_2, \dots, d_n)$ be the diagonal matrix associated to G , where d_i is the degree of vertex v_i . The matrices $L(G) = D(G) - A(G)$ and $Q(G) = D(G) + A(G)$ are respectively called Laplacian and signless Laplacian matrices and their spectrum are respectively called Laplacian spectrum (L -spectrum) and signless Laplacian spectrum (Q -spectrum) of the graph G . Being real symmetric, positive semi-definite matrices, we let $0 = \mu_n \leq \mu_{n-1} \leq \dots \leq \mu_1$ and $0 \leq q_n \leq q_{n-1} \leq \dots \leq q_1$ be the L -spectrum and Q -spectrum of G , respectively. It is well known that $\mu_n = 0$ with multiplicity equal to the number of connected components of G and $\mu_{n-1} > 0$ if and only if G is connected [9]. Moreover, for all $i = 1, 2, \dots, n$, $\mu_i = q_i$ if and only if G is bipartite. For notations and definitions in graphs, we refer to [23].

MSC(2010): Primary: 05C15; Secondary: 05C30; 05C50.

Keywords: Spectra of graph, energy of graph, Laplacian energy, incidence energy

Received: 4 December 2014, Accepted: 29 December 2014.

*Corresponding author.

Let $I(G)$ be the (vertex-edge) incidence matrix of the graph G . For a graph G with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ and edge set $E(G) = \{e_1, e_2, \dots, e_m\}$, the (i, j) -entry of $I(G)$ is 1 if e_j is incident with v_i , and 0 otherwise. As given by Jooyandeh et al. [17] the incidence energy of the incidence matrix $I(G)$ of the graph G is defined as

$$IE = IE(G) = \sum_{k=1}^n \sqrt{\sigma_k},$$

where $\sigma_1, \sigma_2, \dots, \sigma_n$ are the singular values of $I(G)$. Recall that the singular values of a (real) matrix M are equal to the positive square roots of the eigenvalues of MM^t . Various properties and bounds for the incidence energy were recently established in [2, 12, 13, 17, 19, 24, 26, 27, 28]. As is well-known in spectral graph theory that $I(G)I(G)^t = Q(G)$, we can define incidence energy as

$$IE = IE(G) = \sum_{k=1}^n \sqrt{q_k},$$

where q_1, q_2, \dots, q_n are the eigenvalues of $Q(G)$.

Further, Laplacian-spectrum-based graph invariant was put forward by Liu and Liu [20] as

$$LEL = LEL(G) = \sum_{k=1}^{n-1} \sqrt{\mu_k}$$

and was named Laplacian-energy-like invariant. The motivation for introducing LEL was in its analogy to the earlier studied graph energy by Gutman [11] and Laplacian energy by Gutman and Zhou [14]; for more details we refer to [3, 15, 16] and the references cited therein. Recently, several mathematical investigations of LEL were communicated [12, 21, 29]. So several researchers established many lower and upper bounds to estimate the invariant for some classes of graphs [25]. In this paper, we obtain sharp lower bounds for the incidence energy and Laplacian-energy-like invariant which improves some previously known lower bounds for some cases (especially in case of trees, unicyclic, bicyclic, tricyclic, tetracyclic graphs). We also obtain upper bounds for these invariants.

2. Bounds on Laplacian-energy-like invariant

In this section, we obtain upper and lower bounds for the Laplacian-energy-like invariant of a graph. First we start with the following observation by Das [8].

Lemma 2.1. Let G be a graph on $n > 3$ vertices whose distinct Laplacian eigenvalues are $0 < \alpha < \beta$. Then the following hold.

- (i) The multiplicity of α is $n - 2$ if and only if G is one of the graphs $K_{\frac{n}{2}, \frac{n}{2}}$ or $K_{n-1, 1}$.
- (ii) The multiplicity of β is $n - 2$ if and only if G is the graph $K_n - e$.

The following Lemma can be found in [9, 10].

Lemma 2.2. Let G be a connected graph of order n and let Δ be its maximum degree. Then $\Delta + 1 \leq \mu_1 \leq n$. Equality holds on the left if $\Delta = n - 1$ and on the right if and only if G is the join of two graphs.

The following observation can be seen in [18].

Lemma 2.3. Let $G \neq K_n$ be a connected graph of order n and let δ be the smallest vertex degree of G . Then $\mu_{n-1} \leq \delta$, with equality if and only if G is a join of a graph on $\delta(G)$ vertices with another graph.

The next result can be found in [7].

Lemma 2.4.(Pólya-Szegő inequality) Let (a_1, a_2, \dots, a_n) and (b_1, b_2, \dots, b_n) be two sequences of positive real numbers such that there exist positive numbers A, a, B, b satisfying

$$0 < a \leq a_i \leq A < \infty, \quad 0 < b \leq b_i \leq B < \infty,$$

for all $i = 1, 2, \dots, n$. Then

$$(2.1) \quad \frac{\sum_{i=1}^n a_i^2 \sum_{i=1}^n b_i^2}{(\sum_{i=1}^n a_i b_i)^2} \leq \frac{(ab + AB)^2}{4abAB}.$$

The equality holds in (2.1) if and only if $p = \frac{n \cdot \frac{A}{a}}{(\frac{A}{a} + \frac{B}{b})}$, $q = \frac{n \cdot \frac{B}{b}}{(\frac{A}{a} + \frac{B}{b})}$ are integers and if p of the numbers a_1, a_2, \dots, a_n are equal to a and q of these numbers are equal to A , and if the corresponding numbers b_i are equal to b and B , respectively.

We first mention some well known lower bounds for LEL . Gutman et al. [13] obtained the following lower bound.

$$(2.2) \quad LEL(G) \geq \frac{2m}{\sqrt{n}},$$

with equality if and only if $G \cong \overline{K_n}$ or $G \cong K_n$.

Wang et al. [25] obtained the following lower bound.

$$(2.3) \quad LEL(G) \geq \sqrt{\frac{(2m)^3}{2m + n\Delta^2}},$$

where Δ is the maximum vertex degree. Equality occurs if and only if $G \cong K_n$.

It is shown in [25] that the lower bounds (2.2) and (2.3) are incomparable.

We now give a lower bound for LEL in terms of n, m, k .

Theorem 2.5. Let G be a connected graph with n vertices and m edges having algebraic connectivity $\mu_{n-1} \geq k$. Then

$$(2.4) \quad LEL(G) \geq \sqrt{\frac{8m(n-1)\sqrt{kn}}{(\sqrt{n} + \sqrt{k})^2}},$$

with equality if and only if $G \cong K_n$.

Proof. Setting in (2.1) $n = n - 1$, $a_i = \sqrt{\mu_i}$, $b_i = 1$, for $i = 1, 2, \dots, n - 1$ and $a = \sqrt{\mu_{n-1}}$, $A = \sqrt{\mu_1}$, $b = 1$, $B = 1$, we get

$$\frac{\sum_{i=1}^{n-1} \mu_i \sum_{i=1}^{n-1} 1}{(\sum_{i=1}^{n-1} \sqrt{\mu_i})^2} \leq \frac{(\sqrt{\mu_{n-1}} + \sqrt{\mu_1})^2}{4\sqrt{\mu_1\mu_{n-1}}}.$$

This gives,

$$LEL(G) \geq \sqrt{\frac{8m(n-1)\sqrt{\mu_1\mu_{n-1}}}{(\sqrt{\mu_1} + \sqrt{\mu_{n-1}})^2}}.$$

Since,

$$\sqrt{\frac{8m(n-1)\sqrt{\mu_1\mu_{n-1}}}{(\sqrt{\mu_1} + \sqrt{\mu_{n-1}})^2}} \geq \sqrt{\frac{8m(n-1)\sqrt{k\mu_1}}{(\sqrt{\mu_1} + \sqrt{k})^2}},$$

it follows that

$$LEL(G) \geq \sqrt{\frac{8m(n-1)\sqrt{k\mu_1}}{(\sqrt{\mu_1} + \sqrt{k})^2}}.$$

For $x \leq n$, consider the function $f(x) = \frac{8m(n-1)\sqrt{kx}}{(\sqrt{x} + \sqrt{k})^2}$.

For this function, we have

$$f'(x) = \frac{4m(n-1)\sqrt{k}(\sqrt{k} - \sqrt{x})}{\sqrt{x}(\sqrt{x} + \sqrt{k})^3} \leq 0.$$

That is, $f(x)$ is a decreasing function for $x \leq n$. So

$$f(x) \geq f(n) = \frac{8m(n-1)\sqrt{kn}}{(\sqrt{n} + \sqrt{k})^2}.$$

This gives,

$$LEL(G) \geq \sqrt{\frac{8m(n-1)\sqrt{kn}}{(\sqrt{n} + \sqrt{k})^2}}.$$

Equality occurs in (2.4) if and only if equality occurs in (2.1) and $\mu_1 = n$. That is, by Lemma 2.2 and Lemma 2.4, if and only if G is a join of two graphs and p, q are integers, where $p + q = n - 1$ with p of the numbers in $\mu_1, \mu_2, \dots, \mu_{n-1}$ equal to μ_1 and q of them equal to μ_{n-1} . For p, q integers there are $n - 1$ solutions of the equation $p + q = n - 1$ and for any of these integral solutions it follows from Lemma 2.4, that equality occurs if and only if G has two distinct Laplacian eigenvalues. That is, if and only if $G \cong K_n$ [4].

Conversely if $G \cong K_n$, then it is easy to see that equality holds in (2.4). □

Remark 2.6. (i). Let T be a tree of order n , $n \geq 6$ with algebraic connectivity $\mu_{n-1} \geq 0.07$. Then we will show that our lower bound in (2.4) is better than the lower bound in (2.2) for the tree T . For this we have to show that

$$\sqrt{\frac{8m(n-1)\sqrt{(0.07)n}}{(\sqrt{n} + \sqrt{0.07})^2}} \geq \frac{2m}{\sqrt{n}},$$

that is,

$$\frac{80(n-1)^2\sqrt{7n}}{(10\sqrt{n} + \sqrt{7})^2} \geq \frac{4(n-1)^2}{n}, \quad \text{as } m = n - 1$$

that is,

$$20n\sqrt{7n} \geq (10\sqrt{n} + \sqrt{7})^2,$$

which is true for $n \geq 6$. Since for almost all trees algebraic connectivity $\mu_{n-1} \geq 0.07$, it follows that bound (2.4) is better than bound (2.2) for almost all trees.

(ii). Let G be graph of order n having $m \leq \frac{2n(n-1)\sqrt{n}}{(\sqrt{n}+1)^2}$ edges and algebraic connectivity $\mu_{n-1} \geq 1$. Then the lower bound (2.4) is better than the lower bound in (2.2) for G . We have

$$\sqrt{\frac{8m(n-1)\sqrt{kn}}{(\sqrt{n} + \sqrt{k})^2}} \geq \frac{2m}{\sqrt{n}},$$

that is,

$$\frac{8m(n-1)\sqrt{n}}{(\sqrt{n} + 1)^2} \geq \frac{4m^2}{n},$$

that is,

$$(2.5) \quad 2n(n-1)\sqrt{n} \geq m(\sqrt{n} + 1)^2,$$

which is true. In particular if G is unicyclic, bicyclic, tricyclic, tetracyclic graph, then $m = n, n + 1, n + 2, n + 3$ (respectively). It is easy to see that (2.5) holds for $n \geq 5$.

Remark 2.7. (i). Let T be a tree of order n , $n \geq 3$ with maximum degree $\Delta \geq \frac{n}{2}$ and algebraic connectivity $\mu_{n-1} \geq 0.07$. Then the lower bound (2.4) is better than the lower bound (2.3) for T . We have

$$\sqrt{\frac{8m(n-1)\sqrt{kn}}{(\sqrt{n} + \sqrt{k})^2}} \geq \sqrt{\frac{(2m)^3}{2m + n\Delta^2}}$$

that is,

$$\frac{80(n-1)^2\sqrt{7n}}{(10\sqrt{n} + \sqrt{7})^2} \geq \frac{8(n-1)^3}{2(n-1) + n\Delta^2}, \quad \text{as } m = n - 1$$

that is,

$$n\Delta^2 \geq \frac{(n-1)(10\sqrt{n} + \sqrt{7})^2}{10\sqrt{7n}} - 2(n-1)$$

which is true for $\Delta \geq \frac{n}{2}, n \geq 3$.

(ii). Let G be a graph of order n with maximum degree $\Delta \geq \sqrt{\frac{m^2(\sqrt{n}+1)^2}{n(n-1)\sqrt{n}} - \frac{2m}{n}}$ and algebraic connectivity $\mu_{n-1} \geq 1$. Then the bound (2.4) is better than the bound (2.3) for G . We have

$$\frac{8m(n-1)\sqrt{n}}{(\sqrt{n}+1)^2} \geq \frac{8m^3}{2m+n\Delta^2}$$

that is,

$$(2.6) \quad n\Delta^2 \geq \frac{m^2(\sqrt{n}+1)^2}{(n-1)\sqrt{n}} - 2m$$

which is true for $\Delta \geq \sqrt{\frac{m^2(\sqrt{n}+1)^2}{n(n-1)\sqrt{n}} - \frac{2m}{n}}$. In particular if G is unicyclic, bicyclic, tricyclic, then $m = n, n+1, n+2$. It is easy to see that (2.6) holds for $\Delta \geq \frac{n}{2}, n \geq 6$.

We now obtain an upper bound for LEL in terms n, m, Δ, k .

Theorem 2.8. Let G be a connected graph with n vertices and m edges having maximum degree Δ and algebraic connectivity $\mu_{n-1} \geq k$. Then

$$(2.7) \quad LEL(G) \leq \frac{2\sqrt{nk} + k(n-2) + 2m - (\Delta + 1)}{2\sqrt{k}},$$

with equality if and only if $k = n$ and $G \cong K_n$ or $k = 1$ and $G \cong K_{n-1,1}$.

Proof. Let $0 = \mu_n < \mu_{n-1} \leq \mu_{n-2} \leq \dots \leq \mu_2 \leq \mu_1$ be the Laplacian spectrum of G with $\mu_{n-1} \geq k$. Since $\Delta + 1 \leq \mu_1 \leq n$, we have

$$\begin{aligned} LEL(G) &= \sum_{i=1}^{n-1} \sqrt{\mu_i} = \sqrt{\mu_1} + (n-2)\sqrt{\mu_{n-1}} + \sum_{i=2}^{n-2} \left(\frac{\mu_i - \mu_{n-1}}{\sqrt{\mu_i} + \sqrt{\mu_{n-1}}} \right) \\ &\leq \sqrt{\mu_1} + (n-2)\sqrt{\mu_{n-1}} + \sum_{i=2}^{n-2} \left(\frac{\mu_i - \mu_{n-1}}{2\sqrt{\mu_{n-1}}} \right) \\ &= \frac{2\sqrt{\mu_1\mu_{n-1}} + (n-2)\mu_{n-1} + 2m - \mu_1}{2\sqrt{\mu_{n-1}}} \\ &\leq \frac{2\sqrt{n\mu_{n-1}} + (n-2)\mu_{n-1} + 2m - (\Delta + 1)}{2\sqrt{\mu_{n-1}}}. \end{aligned}$$

For $x \geq k$, consider the function

$$f(x) = \frac{2\sqrt{nx} + (n-2)x + 2m - (\Delta + 1)}{2\sqrt{x}}.$$

For this function, we have

$$f'(x) = \frac{x(n-2) + \Delta + 1 - 2m}{4x\sqrt{x}},$$

for all $x \geq k$. Since $(\Delta + 1) + (n - 2)\mu_{n-1} \leq \mu_1 + \mu_2 + \dots + \mu_{n-1} = 2m$, it follows that $f(x)$ is a decreasing function for $x \geq k$. So

$$f(x) \leq f(k) = \frac{2\sqrt{nk} + (n - 2)k + 2m - (\Delta + 1)}{2\sqrt{k}},$$

which gives

$$LEL(G) \leq \frac{2\sqrt{nk} + (n - 2)k + 2m - (\Delta + 1)}{2\sqrt{k}}.$$

Equality occurs in (2.7) if and only if $n = \mu_1 = \Delta + 1$, $\mu_2 = \mu_3 = \dots = \mu_{n-2} = \mu_{n-1} = k$. That is by Lemma 2.2, G is a join of two graphs having at most three distinct Laplacian eigenvalues with $\Delta + 1 = \mu_1$. If G has one distinct Laplacian eigenvalue, then $G \cong \overline{K_n}$, which is not possible as G is connected. If G is a join of two graphs with $\Delta + 1 = \mu_1$ having two distinct Laplacian eigenvalues, then $G \cong K_n$ [4]. If G is join of two graphs having three distinct Laplacian eigenvalues with $\Delta + 1 = \mu_1$, then by Lemma 2.1, $G \cong K_{n-1,1}$.

Conversely if G is one of these graphs, then it is easy to see that equality occurs in (2.7). □

3. Bounds for incidence energy

In order to obtain the bounds for the incidence energy, we need the following Lemmas. Lemma 3.1 can be found in [5].

Lemma 3.1. Let G be a graph of order n having maximum degree Δ , minimum degree δ and largest Q -eigenvalue q_1 . Then $2\delta \leq q_1 \leq 2\Delta$. For a connected graph G , equality holds in either of these inequalities if and only if G is regular.

The next observation is a well known fact [6].

Lemma 3.2. A graph G has two distinct signless Laplacian (Laplacian) eigenvalues if and only if $G \cong K_n$.

The following observation can be found in [5].

Lemma 3.3. For $i = 1, 2, \dots, n$, let μ_i and q_i be the L -spectrum and Q -spectrum of the graph G . Then $\mu_i = q_i$, for all $i = 1, 2, \dots, n$ if and only if G is bipartite.

By Lemma 3.3, it is clear that $IE(G) = LEL(G)$, if and only if G is bipartite. So, if G is bipartite, then any result that holds for $LEL(G)$ also holds for $IE(G)$. Therefore in this section, we consider non-bipartite graphs only.

We first obtain a lower bound for IE in terms of n, m, Δ, k .

Theorem 3.4. Let G be a connected non-bipartite graph with $n > 2$ vertices and m edges having smallest Q -eigenvalue $q_n \geq k$. Then

$$(3.1) \quad IE(G) \geq \sqrt{\frac{8mn\sqrt{2k\Delta}}{(\sqrt{2\Delta} + \sqrt{k})^2}}.$$

Proof. Setting in equation (2.1) $a_i = \sqrt{q_i}$, $b_i = 1$, for $i = 1, 2, \dots, n$ and $a = \sqrt{q_n}$, $A = \sqrt{q_1}$, $b = 1$, $B = 1$, we get

$$\frac{\sum_{i=1}^n q_i \sum_{i=1}^n 1}{(\sum_{i=1}^n \sqrt{q_i})^2} \leq \frac{(\sqrt{q_n} + \sqrt{q_1})^2}{4\sqrt{q_1 q_n}}.$$

This gives,

$$IE(G) \geq \sqrt{\frac{8mn\sqrt{q_1 q_n}}{(\sqrt{q_1} + \sqrt{q_n})^2}}.$$

Since,

$$\sqrt{\frac{8mn\sqrt{q_1 q_n}}{(\sqrt{q_1} + \sqrt{q_n})^2}} \geq \sqrt{\frac{8mn\sqrt{k q_1}}{(\sqrt{q_1} + \sqrt{k})^2}},$$

it follows that

$$IE(G) \geq \sqrt{\frac{8mn\sqrt{k q_1}}{(\sqrt{q_1} + \sqrt{k})^2}}.$$

For $x \leq 2\Delta$, consider the function

$$f(x) = \frac{8mn\sqrt{kx}}{(\sqrt{x} + \sqrt{k})^2}.$$

For this function, we have

$$f'(x) = \frac{4mn\sqrt{k}(\sqrt{k} - \sqrt{x})}{\sqrt{x}(\sqrt{x} + \sqrt{k})^3} \leq 0.$$

That is $f(x)$ is a decreasing function for $x \leq 2\Delta$. So

$$f(x) \geq f(2\Delta) = \frac{8mn\sqrt{2k\Delta}}{(\sqrt{2\Delta} + \sqrt{k})^2}.$$

This gives

$$IE(G) \geq \sqrt{\frac{8mn\sqrt{2k\Delta}}{(\sqrt{2\Delta} + \sqrt{k})^2}}.$$

Equality occurs in (3.1) if and only if equality occurs in (2.1), $q_1 = 2\Delta$ and $q_n = k$. That is, by Lemma 2.2 and Lemma 3.1, if and only if G is regular and p, q are integers, where $p + q = n$ with p of the numbers in q_1, q_2, \dots, q_n equal to $A = q_1$ and q of them equal to $a = q_n$. For p, q integers there are n solutions of the equation $p + q = n$ and for any of these integral solutions it follows from Lemma 2.4, that equality occurs if and only if G has two distinct Q -eigenvalues. By Lemma 3.2, a graph with two distinct Q -eigenvalues is K_n , having $A = q_1 = 2(n - 1)$, $p = 1$ and $a = q_n = n - 2$, $q = n - 1$. For

this choice of A, a, p, q , we have from Lemma 2.2, $1 = \frac{n\sqrt{\frac{2(n-1)}{n-2}}}{1+\sqrt{\frac{2(n-1)}{n-2}}}$. That is, $2(n-2)(n-1) = 1$, which gives, $n = \frac{3 \pm \sqrt{3}}{2}$, which is not possible. So equality does not occur in (3.1) for any graph G . \square

Since $q_1 \leq 2\Delta \leq 2(n-1)$, it follows from Theorem 3.4,

$$f(q_1) \geq f(2(n-1)) = \sqrt{\frac{8mn\sqrt{2k(n-1)}}{(\sqrt{2(n-1)} + \sqrt{k})^2}}.$$

Therefore we have the following observation.

Corollary 3.5. Let G be a connected non-bipartite graph with n vertices and m edges having smallest Q -eigenvalue $q_n \geq k$. Then

$$(3.2) \quad IE(G) \geq \sqrt{\frac{8mn\sqrt{2k(n-1)}}{(\sqrt{2(n-1)} + \sqrt{k})^2}}.$$

Recall from [13] that a lower bound for IE was given as follows.

Lemma 3.6. Let G be graph with n vertices and m edges. Then

$$(3.3) \quad IE(G) \geq \frac{2m}{\sqrt{n}},$$

with equality if and only if $G \cong \overline{K_n}$ or $G \cong K_2$.

Remark 3.7. Let G be a non-bipartite graph with n vertices, smallest Q -eigenvalue $q_n \geq k = 0.5$ and $m \leq \frac{2n^2\sqrt{n-1}}{(\sqrt{2n-2} + \sqrt{0.5})^2}$. Then the bound (3.2) is better than the bound (3.3) for G . For this we need to show that

$$\frac{\sqrt{8mn\sqrt{2(0.5)(n-1)}}}{\sqrt{2(n-1)} + \sqrt{0.5}} \geq \frac{2m}{\sqrt{n}},$$

that is,

$$(3.4) \quad 2n^2\sqrt{n-1} \geq m(\sqrt{2n-2} + \sqrt{0.5})^2,$$

which is true. In particular if G is a unicyclic, bicyclic, tricyclic, tetracyclic graph, then $m = n, n+1, n+2, n+3$ (respectively). It is easy to see that (3.4) holds for $n \geq 5$.

We now have an upper bound for IE in terms of n, m, Δ, δ, k .

Theorem 3.8. Let G be a non-bipartite connected graph with n vertices, m edges having maximum degree Δ , minimum degree δ and smallest Q -eigenvalue $q_n \geq k$. Then

$$IE(G) \leq \frac{2\sqrt{2k\Delta} + (n-1)k + 2m - 2\delta}{2\sqrt{k}},$$

with equality if and only if $G \cong K_n$.

Proof. This follows by proceeding similarly as in Theorem 2.4.

REFERENCES

- [1] F. Ayoobi, G. R. Omid and B. Tayfeh-Rezaie, A note on graphs whose signless Laplacian has three distinct eigenvalues, *Linear and Multilinear Algebra*, **59** (2011) 701–706.
- [2] S. B. Bozkurt and I. Gutman, Estimating the incidence energy, *MATCH Commun. Math. Comput. Chem.*, **70** (2013) 143–156.
- [3] T. A. Chishti, Hilal A. Ganie and S. Pirzada, Properties of strong double graphs, *J. Discrete Math. Sci. Cryptogr.*, **17** (2014) 311–319.
- [4] D. Cvetković, M. Doob and H. Sachs, Spectra of graphs—Theory and Application, Academic Press, New York, 1980.
- [5] D. Cvetković, P. Rowlinson and S. K. Simić, Signless Laplacians of finite graphs, *Linear Algebra Appl.*, **423** (2007) 155–171.
- [6] D. Cvetković and S. K. Simić, Towards a spectral theory of graphs based on the signless Laplacian, I, *Publ. Inst. Math.*, (Beograd) (N. S.), **85** (2009) 19–33.
- [7] S. S. Dragomir, A survey on Cauchy-Bunyakovsky-Schwarz type discrete inequalities, *JIPAM. J. Inequal. Pure Appl. Math.*, Article 63, **4** (2003) pp. 142.
- [8] K. C. Das, A sharp upper bound for the number of spanning trees of a graph, *Graphs Combin.*, **23** (2007) 625–632.
- [9] M. Fiedler, Algebraic Connectivity of Graphs, *Czechoslovak Math. J.*, **23** (1973) 298–305.
- [10] R. Grone and R. Merris, The Laplacian spectrum of a graph II, *SIAM J. Discrete Math.*, **7** (1994) 221–229.
- [11] I. Gutman, The energy of a graph: old and new results, Algebraic combinatorics and applications (Gweinstein, 1999), Springer, Berlin, 2001 196–211.
- [12] I. Gutman, D. Kiani and M. Mirzakhah, On incidence energy of graphs, *MATCH Commun. Math. Comput. Chem.*, **62** (2009) 573–580.
- [13] I. Gutman, D. Kiani, M. Mirzakhah and B. Zhou, On incidence energy of a graph, *Linear Algebra Appl.*, **431** (2009) 1223–1233.
- [14] I. Gutman and B. Zhou, Laplacian energy of a graph, *Linear Algebra Appl.*, **414** (2006) 29–37.
- [15] I. Gutman, B. Zhou and B. Furtula, The Laplacian-energy like invariant is an energy like invariant, *MATCH Commun. Math. Comput. Chem.*, **64** (2010) 85–96.
- [16] H. A. Ganie, S. Pirzada and I. Antal, Energy and Laplacian energy of double graphs, *Acta Univ. Sap. Informatica*, **6** (2014) 89–116.
- [17] M. Jooyandeh, D. Kiani and M. Mirzakhah, Incidence energy of a graph, *MATCH Commun. Math. Comput. Chem.*, **62** (2009) 561–572.
- [18] S. J. Kirkland, J. J. Moliterno, M. Neumann and B. L. Shader, On graphs with equal algebraic and vertex connectivity, *Linear Algebra Appl.*, **341** (2002) 45–56.
- [19] M. Liu and B. Liu, On sum of powers of the signless Laplacian eigenvalues of graphs, *Hacet. J. Math. Stat.*, **41** (2012) 527–536.
- [20] J. Liu and B. Liu, A Laplacian-energy-like invariant of a graph, *MATCH Commun. Math. Comput. Chem.*, **59** (2008) 355–372.
- [21] M. Liu, B. Liu and X. Tan, The first to ninth greatest LEL-invariants of connected graphs, *Util. Math.*, **93** (2014) 153–160.
- [22] A. Mohammadian and B. Tayfeh-Rezaie, Graphs with four distinct Laplacian eigenvalues, *J. Algebraic Combin.*, **34** (2011) 671–682.

- [23] S. Pirzada, An Introduction to Graph Theory, Universities Press, Orient Blackswan, 2012.
- [24] Z. Tang and Y. Hou, On incidence energy of trees, *MATCH Commun. Math. Comput. Chem.*, **66** (2011) 977–984.
- [25] W. Wang and Y. Luo, On Laplacian-energy-like invariant of a graph, *Linear Algebra Appl.*, **437** (2012) 713–721.
- [26] B. Zhou, More upper bounds for the incidence energy, *MATCH Commun. Math. Comput. Chem.*, **64** (2010) 123–128.
- [27] J. Zhang and J. Li, New results on the incidence energy of graphs, *MATCH Commun. Math. Comput. Chem.*, **68** (2012) 777-803.
- [28] B. Zhou and A. Ilić, On the sum of powers of Laplacian eigenvalues of bipartite graphs, *Czechoslovak Math. J.*, **60** (2010) 1161–1169.
- [29] B. X. Zhu, The Laplacian-energy like of graphs, *Appl. Math. Lett.*, **24** (2011) 1604–1607.

S. Pirzada

Department of Mathematics, University of Kashmir, Srinagar, India

Email: pirzadasd@kashmiruniversity.ac.in

Hilal A. Ganie

Department of Mathematics, University of Kashmir, Srinagar, India

Email: hilahmad1119kt@gmail.com