



www.combinatorics.ir

Transactions on Combinatorics

ISSN (print): 2251-8657, ISSN (on-line): 2251-8665

Vol. 01 No.1 (2012), pp. 45-51.

© 2012 University of Isfahan



www.ui.ac.ir

ON THE TOTAL DOMATIC NUMBER OF REGULAR GRAPHS

H. ARAM, S. M. SHEIKHOESLAMI* AND L. VOLKMANN

Communicated by Manoucher Zaker

ABSTRACT. A set S of vertices of a graph $G = (V, E)$ without isolated vertex is a *total dominating set* if every vertex of $V(G)$ is adjacent to some vertex in S . The *total domatic number* of a graph G is the maximum number of total dominating sets into which the vertex set of G can be partitioned. We show that the total domatic number of a random r -regular graph is almost surely at most $r - 1$, and that for 3-regular random graphs, the total domatic number is almost surely equal to 2. We also give a lower bound on the total domatic number of a graph in terms of order, minimum degree and maximum degree. As a corollary, we obtain the result that the total domatic number of an r -regular graph is at least $r/(3\ln(r))$.

1. Introduction

Let $G = (V(G), E(G)) = (V, E)$ be a simple graph of order n with minimum degree $\delta(G) \geq 1$. The neighborhood of a vertex u is denoted by $N_G(u)$ and its degree $|N_G(u)|$ by $d_G(u)$ (briefly $N(u)$ and $d(u)$ when no ambiguity on the graph is possible). The *minimum* and *maximum degree* of a graph G are denoted by $\delta = \delta(G)$ and $\Delta = \Delta(G)$, respectively. The *open neighborhood* of a set $S \subseteq V$ is the set $N(S) = \cup_{v \in S} N(v)$, and the *closed neighborhood* of S is the set $N[S] = N(S) \cup S$. A *matching* is a set of edges with no shared endvertices. A perfect matching M of G is a matching with $V(M) = V(G)$. The maximum number of edges of a matching in G is denoted by $\alpha'(G)$ (α' for short). If $C = (v_1, v_2, \dots, v_n)$ is a cycle and v_i, v_k are distinct vertices of C , then the segment $[v_i, v_k]$ of C is defined as the set $\{v_i, v_{i+1}, v_{i+2}, \dots, v_k\}$, where the subscripts are taken modulo n . If $f(n)$ and $g(n)$ are real valued functions of an integer variable n , then we write $f(n) = O(g(n))$ (or $f(n) = \Omega(g(n))$)

MSC(2010): 05C69.

Keywords: total dominating set, total domination number, total domatic number, regular graph.

Received: 12 January 2012, Accepted: 19 March 2012.

*Corresponding author.

if there exist constants $C > 0$ and n_0 such that $f(n) \leq Cg(n)$ (or $f(n) \geq Cg(n)$) for $n \geq n_0$. We also write $f(n) \sim g(n)$ if $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$. We use [9] for terminology and notation which are not defined here.

A set S of vertices of a graph G with minimum degree $\delta(G) > 0$ is a *total dominating set* if $N(S) = V(G)$. The minimum cardinality of a total dominating set, denoted by $\gamma_t(G)$, is called the *total domination number* of G . A $\gamma_t(G)$ -*set* is a total dominating set of G of cardinality $\gamma_t(G)$.

A partition of $V(G)$, all of whose classes are total dominating sets in G , is called a *total domatic partition* of G . The maximum number of classes of a total domatic partition of G is called the *total domatic number* of G and is denoted by $d_t(G)$. The total domatic number was introduced by Cockayne, Dawes and Hedetniemi in [5] and has been studied by several authors (see for example, [2, 4, 11, 12]). More information on the total domination number and the total domatic number can be found in the monographs [7, 8] by Haynes, Hedetniemi and Slater.

We use the following standard model $\mathcal{G}_{n,r}$ to generate r -regular graphs on n vertices uniformly: to construct a random r -regular graph on the vertex set $\{v_1, v_2, \dots, v_n\}$, take a random matching on the vertex set $\{v_{1,1}, v_{1,2}, \dots, v_{1,r}, v_{2,1}, \dots, v_{2,r}, \dots, v_{n,r}\}$ and collapse each set $\{v_{i,1}, v_{i,2}, \dots, v_{i,r}\}$ into a single vertex v_i . If the resulting graph contains any loops or multiple edges, discard it. All r -regular graphs are generated uniformly with this method. Wormald [10] has shown that 3-regular graphs are almost surely Hamiltonian, and that the model $\mathcal{G}_{n,r}$ and $\mathcal{H}_n \oplus \mathcal{G}_{n,r-2}$ are contiguous, meaning roughly that events that are almost sure in one model are almost sure in the other. Thus if an event is almost surely true in a random graph constructed from a random Hamilton cycle plus a random matching, then it is almost surely true in a random 3-regular graph. For more details the reader is referred to [10].

We make use of the following results.

Theorem A. ([10]) If G is a 3-regular random graph, then a.a. G consists of a Hamilton cycle plus a random matching.

Theorem B. ([5]) For every graph G of order n without isolated vertices,

$$d_t(G) \leq \min \left\{ \delta(G), \frac{n}{\gamma_t(G)} \right\}.$$

2. A lower bound on the total domatic number

In this section we will show that the total domatic number of a random 3-regular graph is at least 2.

Definition 2.1. Let G be a 3-regular graph obtained from a cycle $C = (v_1, v_2, \dots, v_n)$ by adding a perfect matching M . An edge $v_i v_{i+1}$ of C (the indices are taken modulo n) is a 4-edge if v_i and v_{i+1} have matching partners v_j and v_k respectively, such that the cycle segments $[v_j, v_i]$ and $[v_{i+1}, v_k]$ are disjoint and have cardinality 0 (mod 4).

Lemma 2.2. Let $G = C \cup M$ as above. If C has a 4-edge then $d_t(G) \geq 2$.

Proof. Let $v_i v_{i+1}$ be a 4-edge of C , and let v_j, v_k be their matching partners, respectively. Without loss of generality, we may assume that $i = 1$. If $n \equiv 0 \pmod{4}$, then obviously $S_1 = \{v_{4i+1}, v_{4i+2} \mid 0 \leq i \leq \frac{n}{4} - 1\}$ and $S_2 = V(G) - S_1$ are two disjoint total dominating sets and hence $d_t(G) \geq 2$.

Now let $n \equiv 2 \pmod{4}$. Then $n = 4s + 2$ for some positive integer s . If $n = 6$, then the result is immediate. Assume that $n \geq 10$. Then $k \equiv 1 \pmod{4}$ and $j \equiv 0 \pmod{4}$. Let $k = 4\ell + 1$, $j = 4r$ and define $S_1 = \{v_{4i+1}, v_{4i+2} \mid 0 \leq i \leq \frac{k-1}{4} - 1\} \cup \{v_{4i+3}, v_{4i} \mid r \leq i \leq s\}$ if $j = k + 3$, or $S_1 = \{v_{4i+1}, v_{4i+2} \mid 0 \leq i \leq \frac{k-1}{4} - 1\} \cup \{v_{4i+3}, v_{4i} \mid r \leq i \leq s\} \cup \{v_{4i}, v_{4i+1} \mid \ell + 1 \leq i \leq r - 1\}$ when $j > k + 3$ and $S_2 = V(G) - S_1$. Obviously, S_1 and S_2 are two disjoint total dominating sets. Thus $d_t(G) \geq 2$ and the proof is complete. \square

The proof of the following lemma is essentially similar to the proof of Lemma 2 of [6].

Lemma 2.3. *Let G be a graph obtained from a cycle $C = (v_1, v_2, \dots, v_n)$ of even order by adding a random matching M . Then G has a 4-edge a.a.*

Proof. Define random variables X_i for $i = 1, 2, \dots, n$ by

$$X_i = \begin{cases} 1 & \text{if } v_i v_{i+1} \in E(C) \text{ is a 4-edge} \\ 0 & \text{otherwise.} \end{cases}$$

and let $X = \sum_{i=1}^n X_i$. Then each X_i has expectation

$$E(X_i) = P(X_i = 1) = \frac{1}{2} \left(\frac{1}{4} + \frac{1}{n} \right)^2 = \frac{1}{32} + O(1/n),$$

and variance

$$\text{var}(X_i) = E(X_i^2) - E(X_i)^2 = E(X_i) - E(X_i)^2 = \frac{31}{1024} + O(1/n).$$

The covariance of X_i and X_j for $i < j$ equals

$$\begin{aligned} \text{cov}(X_i, X_j) &= E(X_i X_j) - E(X_i)E(X_j) \\ &= \begin{cases} 1/1024 - (1/32)^2 + O(1/n) & \text{if } i < j - 1 \\ \frac{8}{3} \left(\frac{1}{1024} \right) - (1/32)^2 + O(1/n) & \text{if } i = j - 1 \text{ and } n \equiv 2 \pmod{4} \\ 0 - (1/32)^2 + O(1/n) & \text{if } i = j - 1 \text{ and } n \equiv 0, 1, 3 \pmod{4} \end{cases} \\ &= \begin{cases} O(1/n) & \text{if } i < j - 1 \\ O(1) & \text{if } i = j - 1. \end{cases} \end{aligned}$$

Note that $X_i X_{i+1} = 1$ implies that $n \equiv 2 \pmod{4}$. To see this, let v_k be the matching partner of v_{i+1} . If $X_i X_{i+1} = 1$, then $v_i v_{i+1}$ and $v_{i+1} v_{i+2}$ are 4-edges and thus $n + 2 \equiv |[v_{i+1}, v_k]| + |[v_k, v_{i+1}]| \equiv 0 + 0 \equiv 0 \pmod{4}$, i.e., $n \equiv 2 \pmod{4}$.

Hence the random variable X has expectation

$$E(X) = \sum_{i=1}^n E(X_i) = n/32 + O(1) = O(n)$$

and variance

$$\begin{aligned} \text{var}(X) &= \sum_{i=1}^n \text{var}(X_i) + 2 \sum_{i < j-1} \text{cov}(X_i, X_j) + 2 \sum_{i=1}^n \text{cov}(X_i, X_{i+1}) \\ &= \frac{31}{1024}n + 2 \sum_{i < j-1} O(1/n) + 2 \sum_{i=1}^n O(1) \\ &= O(n) \end{aligned}$$

By Chebyshev's inequality, we have

$$\text{prob}(X = 0) \leq \frac{\text{var}(X)}{E(X)^2} = \frac{O(n)}{(O(n))^2} = O(1/n).$$

Hence $X > 0$ a.a., i.e., G has a 4-edge. □

An immediate consequence of Theorem A and Lemmas 2.2 and 2.3 now follows.

Theorem 2.4. If G is a random 3-regular graph, then $d_t(G) \geq 2$ a.a.

3. An upper bound for the total domatic number

The proof of the following theorem is essentially similar to the proof of Theorem 2 of [6].

Theorem 3.1. Let $r \geq 3$ and let G be a random r -regular graph. Then $d_t(G) \leq r - 1$.

Proof. Suppose to the contrary that G is an r -regular graph with $d_t(G) > r - 1$. It follows from Theorem B that $d_t(G) = r$. Let V_1, V_2, \dots, V_r be a total domatic partition for G . Then each vertex has a neighbor in every V_i . Since every vertex has precisely r neighbors, we deduce that every vertex in V_i has precisely one neighbor in V_j for each j . Hence,

$$|N(v) \cap V_i| = 1 \quad \text{for all } v \in V(G) \text{ and } i \in \{1, 2, \dots, r\}.$$

For $i \neq j$ we deduce that

$$(3.1) \quad E_{ij} := \{uv \in E(G) | u \in V_i, v \in V_j\} \text{ is a perfect } V_i - V_j \text{ matching}$$

and so

$$(3.2) \quad |V_1| = |V_2| = \dots = |V_r| = \frac{n}{r}.$$

It follows from the above argument that every r -regular graph with $d_t = r$ on the vertex set $V(G)$ can be obtained by first partitioning $V(G)$ into r sets, all of equal cardinality, and then adding a perfect matching between the vertices of every partition, implying that n/r is even, and finally adding a perfect matching between all pairs of partition sets. Suppose n is a multiple of r . Since the sets are not distinguishable, the first step can be done in

$$\binom{n}{n/r, n/r, \dots, n/r} \frac{1}{r!}$$

ways, the second step can be done in

$$\left[\left(\frac{n}{r} - 1\right) \left(\frac{n}{r} - 3\right) \dots 1 \right]^r = \frac{\left(\left(\frac{n}{r}\right)!\right)^r}{2^{n/2} \left(\left(\frac{n}{2r}\right)!\right)^r}$$

ways, and the last step can be done in

$$\left(\left(\frac{n}{r}\right)!\right)^{\binom{r}{2}}$$

ways, since there are $\binom{r}{2}$ different pairs of sets V_i, V_j , and between each pair a matching can be added in $\left(\frac{n}{r}\right)!$ ways. Hence, an upper bound on the number of labeled r -regular graphs of order n with $d_t = r$, is

$$\begin{aligned} & \binom{n}{n/r, n/r, \dots, n/r} \frac{1}{r!} \left(\left(\frac{n}{r}\right)!\right)^{\binom{r}{2}} \frac{\left(\left(\frac{n}{r}\right)!\right)^r}{2^{n/2} \left(\left(\frac{n}{2r}\right)!\right)^r} \\ &= \frac{n!}{\left(\left(\frac{n}{r}\right)!\right)^r r!} \left(\left(\frac{n}{r}\right)!\right)^{\frac{r(r-1)}{2}} \frac{\left(\left(\frac{n}{r}\right)!\right)^r}{2^{n/2} \left(\left(\frac{n}{2r}\right)!\right)^r}, \end{aligned}$$

and hence, by Stirling's formula ($n! \sim \left(\frac{n}{e}\right)^n \sqrt{2\pi n} (1 + \frac{1}{12n} + O(\frac{1}{n^2}))$) the upper bound is, for large n and constant r ,

$$\left(\frac{n}{e}\right)^n \sqrt{2\pi n} \left(1 + \frac{1}{12n} + O\left(\frac{1}{n^2}\right)\right) \cdot \frac{1}{r! 2^{n/2}} \cdot \left(\frac{n}{re}\right)^{\frac{r(r-1)}{2}}$$

$$\cdot \left[\sqrt{2\pi n/r} \left(1 + \frac{1}{12n} + O\left(\frac{1}{n^2}\right)\right)\right]^{\frac{r(r-1)}{2}} \cdot \frac{1}{\left(\frac{n}{2re}\right)^{rn} \left(\sqrt{\pi n/r} \left(1 + \frac{1}{12n} + O\left(\frac{1}{n^2}\right)\right)\right)^r}.$$

Denote this last expression by $\text{DOMT}(r, n)$. The total number of r -regular graphs, as given in [3] is asymptotic to

$$e^{-(r^2-1)/4} \frac{(rn)!}{(rn/2)! 2^{rn/2} (r!)^n} = \frac{e^{-(r^2-1)/4}}{2^{rn/2} (r!)^n} \cdot \frac{\left(\frac{rn}{e}\right)^n \sqrt{2\pi rn} \left(1 + \frac{1}{12n} + O\left(\frac{1}{n^2}\right)\right)}{\left(\frac{rn}{2e}\right)^n \sqrt{\pi rn} \left(1 + \frac{1}{12n} + O\left(\frac{1}{n^2}\right)\right)}$$

Denote this last expression by $\text{TOTAL}(r, n)$. Then the proportion of r -regular graphs with $d_t = r$, $\text{DOMT}(r, n)/\text{TOTAL}(r, n)$, is at most

$$\left(\frac{r!}{r^{r-1}}\right)^n O\left(n^{\frac{(r-1)(r-2)}{4}}\right).$$

Since $\frac{r!}{r^{r-1}}$ is less than 1, so the limit $\text{DOMT}(r, n)/\text{TOTAL}(r, n)$ tends to 0, as desired. This completes the proof. \square

4. Total domatic number and minimum degree

If G is a graph of order n , then Zelinka [12] gave the following lower bound on the total domatic number

$$d_t(G) \geq \lfloor \frac{n}{n - \delta(G) + 1} \rfloor.$$

The proof of the following theorem is essentially similar to the proof of Theorem 3 of [6] and we leave it to the reader.

Theorem 4.1. Let G be a graph of order n with minimum degree δ and maximum degree Δ , and let k be a nonnegative integer. If

$$e(\Delta^2 + 1)k \left(1 - \frac{1}{k}\right)^\delta < 1,$$

then $d_t(G) \geq k$.

For the special case of a regular graph, we obtain a significant improvement of Zelinka's bound.

Corollary 4.2. Let G be an r -regular graph with $r \geq 3$. Then

$$d_t(G) \geq \frac{r}{3 \ln r}.$$

Proof. With $\Delta = \delta = r$ and $k = \frac{r}{3 \ln r}$ we have

$$\begin{aligned} e(\Delta^2 + 1)k(1 - \frac{1}{k})^\delta &= e(r^2 + 1)k(1 - \frac{1}{k})^r \\ &\leq e(r^2 + 1)\frac{r}{3 \ln r}e^{-\frac{3 \ln r}{r}r} \\ &= \frac{e(r^2 + 1)}{3r^2 \ln r} \\ &< 1. \end{aligned}$$

Now it follows from Theorem 4.1 that $d_t(G) \geq \frac{r}{3 \ln(r)}$. □

A question that arises naturally is whether the bound in Corollary 4.2 is best possible. For a positive integer r , let $f(r)$ be the minimum total domatic number of all r -regular graphs. By Corollary 4.2 we have $f(r) \geq \frac{r}{3 \ln(r)}$. On the other hand, it follows from [1] that there exist r -regular graphs of order n with total domination number $(1 + o(1))\frac{n \ln(r)}{r}$. According to Theorem B, the total domatic number of those graphs is at most $n/\gamma_t(G) = (1 + o(1))\frac{r}{\ln(r)}$. This proves $f(r) = \Omega(\frac{r}{\ln(r)})$, and the order of magnitude of the bound in Corollary 4.2 is best possible.

REFERENCES

- [1] N. Alon, Transversal numbers of uniform hypergraphs, *Graphs Combin.* **6** (1990), 1–4.
- [2] S. Arumugam and A. Thuraiwamy, Total domatic number of a graph, *Indian J. Pure Appl. Math.* **29** (1998), 513–515.
- [3] E. A. Bender and E. R. Canfield, The asymptotic number of labeled graphs with given degree sequences, *J. Combin. Theory Ser. A* **24** (1978), 296–307.
- [4] I. Bouchemakh and S. Ouatiki, On the domatic and the total domatic numbers of the 2-section graph of the order-interval hypergraph of a finite poset, *Discrete Math.* **309** (2009), 3674–3679.
- [5] E. J. Cockayne, R. M. Dawes, S. T. Hedetniemi, Total domination in graphs, *Networks* **10** (1980), 211–219.
- [6] P. Dankelmann, N. J. Calkin, The Domatic Number of Regular Graphs, *Ars Combin.* **73** (2004), 247–255.
- [7] T. W. Haynes, S. T. Hedetniemi, P. J. Slater, *Fundamentals of Domination in Graphs*, Marcel Dekker, Inc., New York, 1998.
- [8] T. W. Haynes, S. T. Hedetniemi, P. J. Slater, editors, *Domination in Graphs, Advanced Topics*, Marcel Dekker, Inc., New York, 1998.
- [9] D. B. West, *Introduction to Graph Theory*, Prentice-Hall, Inc, 2000.
- [10] N. C. Wormald, *Models of Random Regular Graphs*. In: Surveys in Combinatorics 1999 (J.D. Lamb and D.A. Preece, eds., 239–298), London Mathematical Society Lecture Notes Series 267, Cambridge University Press, Cambridge, 1999.
- [11] B. Zelinka, Total domatic number of cacti, *Math. Slovaca* **38** (1988), 207–214.
- [12] B. Zelinka, Total domatic number and degrees of vertices of a graph, *Math. Slovaca* **39** (1989), 7–11.

H. Aram

Department of Mathematics, Azarbaijan University of Tarbiat Moallem, Tabriz, I.R. Iran

Email: hamideh.aram@gmail.com

S. M. Sheikholeslami

Department of Mathematics, Azarbaijan University of Tarbiat Moallem, Tabriz, I.R. Iran

Email: s.m.sheikholeslami@azaruniv.edu

L. Volkmann

Lehrstuhl II für Mathematik, RWTH-Aachen University, 52056 Aachen, Germany

Email: volkm@math2.rwth-aachen.de